

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/025, C12N 15/37, 15/86, 5/10, A61K 39/12, 31/73	A1	(11) International Publication Number: WO 97/05164 (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/AU96/00473 (22) International Filing Date: 26 July 1996 (26.07.96) (30) Priority Data: PN 4439 27 July 1995 (27.07.95) AU (71) Applicants (for all designated States except US): CSL LIMITED [AU/AU]; 45 Poplar Road, Parkville, VIC 3052 (AU). THE UNIVERSITY OF QUEENSLAND [AU/AU]; St. Lucia, QLD 4067 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only): WEBB, Elizabeth, Ann [AU/AU]; 36 Zig Zag Road, Eltham, VIC 3095 (AU). MARGETTS, Mary, Brigid [IE/AU]; 92 Bent Street, Moonee Ponds, VIC 3039 (AU). COX, John, Cooper [AU/AU]; 365 Bacchus Marsh Road, Bullengarook, VIC 3437 (AU). FRAZER, Ian [AU/AU]; 64 Eighth Avenue, St. Lucia, QLD 4067 (AU). McMILLAN, Nigel, Alan, John [NZ/AU]; Center for Immunology and Cancer Research, Princess Alexandra Hospital, University of Queensland, Wollongabba, QLD 4102 (AU). WILLIAMS, Mark, Philip [AU/AU]; 1/58 Chester Road, Annerley, QLD 4103 (AU). MOLONEY, Margaret, Bridget, Holland [AU/AU]; 22 Leslie Road,	(74) Agents: SLATTERY, John, Michael et al.; Davies Collison Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU). (81) Designated States: AU, BR, CA, IL, JP, KR, NZ, SG, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: PAPILLOMAVIRUS POLYPROTEIN CONSTRUCTS		
(57) Abstract <p>A papillomavirus polyprotein construct comprises at least two amino acid sequences fused directly or indirectly together, each of the sequences being the sequence of an early ORF protein of papillomavirus or an immunogenic variant or fragment thereof, and at least one of said sequences being other than the E6 or E7 protein sequence or an immunogenic variant or fragment thereof. Nucleic acid molecules encoding the polyprotein construct, prophylactic or therapeutic compositions comprising the polyprotein construct or the nucleic acid molecule, and methods for eliciting an immune response against papillomavirus in a host animal are also provided.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

"PAPILLOMAVIRUS POLYPROTEIN CONSTRUCTS"

FIELD OF THE INVENTION

5 This invention relates to polyprotein constructs and in particular polyprotein constructs comprising a plurality of papillomavirus (PV) amino acid sequences which may be used in compositions for eliciting an immune response against PV, and particularly human papillomavirus (HPV), in a host animal.

10 BACKGROUND OF THE INVENTION

Papillomaviruses induce benign hyperproliferative lesions in humans and in many animal species, some of which undergo malignant conversion. The biology of papillomavirus infection is summarised in a review by J.P. Sundberg, entitled
15 "Papillomavirus Infections in Animals" In "Papillomaviruses and Human Disease" edited by K. Syrjanen, L. Gissmann and L.G. Koss, Springer Verlag (1987).

Papillomaviruses are a family of small DNA viruses encoding up to eight early (E1, E2, E3, E4, E5, E6, E7 and E8) and two late genes (L1 and L2). These viruses have been
20 classified in several distinct groups such as HPV which are differentiated into types 1 to ~ 70 depending upon DNA sequence homology. A clinicopathological grouping of HPV and the malignant potential of the lesions with which they are most frequently associated are summarised in "Papillomaviruses and Human Cancer" by H. Pfister, CRC Press, Inc. (1990). For example, HPV type 1 (HPV-1) is present in plantar warts, HPV-6 or HPV-11
25 are associated with condylomata acuminata (anogenital warts), and HPV-16 or HPV-18 are common in pre-malignant and malignant lesions of the cervical squamous epithelium.

The immunological approach to the prevention of HPV disease requires a thorough analysis of the viral proteins against which humoral and cellular immune
30 responses are mounted during and after infection. However, despite recent limited

- 2 -

success (Kreider *et al.*, 1986, *J. Virol.*, **59**, 369; Sterling *et al.*, 1990, *J. Virol.*, **64**, 6305; Meyers *et al.*, 1992, *Science*, **257**, 971; Dollard *et al.*, 1992, *Genes and Development*, **6**, 1131), papillomaviruses are notoriously refractory to growth in cultured cells (Teichman and LaPorta, 1987 *In "The Papovaviridae"*, Vol 2 edited by N.P. Salzman and P.M. Howley, p.109). As a consequence, the lack of viral reagents has delayed the analysis of the immune response to PV infection.

The recent advent of recombinant expression systems *in vitro* has allowed the production of viral proteins encoded by both early and late genes in relatively large amounts and in a purified form (Tindle *et al.*, 1990, *J. Gen. Virol.*, **71**, 1347; Jarrett *et al.*, 1991, *Virology*, **184**, 33; Ghim *et al.*, 1992, *Virology*, **190**, 548; Stacey *et al.*, 1991, *J. Gen. Virol.*, **73**, 2337). These systems have, for the first time, allowed the analysis of the host immune response to these viral proteins.

Interest in immune responses to the non-structural early open reading frame (ORF) proteins of HPV has centred on HPV-16 E7 because of an apparent association between serum antibodies to this protein and cervical cancer (for a review, see "Immune Response to Human Papillomaviruses and the Prospects of Human Papillomavirus-Specific Immunisation" by Tindle and Frazer *In "Human Pathogenic Papillomaviruses"* edited by H. zur Hausen, Current Topics in Microbiology Immunology, **186**, Springer-Verlag, Berlin, 1994).

The immune responses to other HPV early ORF proteins have also been investigated including HPV-16 E6 (Stacey *et al.*, 1992, *J. Gen. Virol.*, **73**, 2337; Bleul *et al.*, 1991, *J. Clin. Microbiol.*, **29**, 1579; Dillner, 1990, *Int. J. Cancer*, **46**, 703; and Müller *et al.*, 1992, *Virology*, **187**, 508), HPV-16 E2 (Dillner *et al.*, 1989 *Proc. Natl. Acad. Sci. USA*, **86**, 3838; Dillner, 1990, *supra*; Lehtinen *et al.*, 1992, *J. Med. Virol.*, **37**, 180; Mann *et al.*, 1990, *Cancer Res.*, **50**, 7815; and Jenison *et al.*, 1990, *J. Infect. Dis.*, **162**, 60) and HPV-16 E4 (Köchel *et al.*, 1991, *Int. J. Cancer*, **48**, 682; Jochmus-Kudielka *et al.*, 1989, *JNCI*, **81**, 1698; and Barber *et al.*, 1992, *Cancer Immunol. Immunother.*, **35**,

33). However, comparison of these studies reveals a lack of correlation between the results of the various assays which have been used in assessing HPV early ORF protein reactivity in serum (Tindle and Frazer, 1994, *supra*).

5 In addition, antibodies to other HPV early ORF proteins have not yet been sought with sufficient rigour in large enough numbers of patients to determine their utility as disease markers or as indicators of HPV protein immunogenicity following HPV infection.

A problem associated with immunising animals with preparations of individual PV
10 proteins is that most of these proteins are comparatively small and might therefore not comprise many reactive epitopes. In addition, immunodominance of particular B or T cell epitopes within a single PV protein would vary presumably between animals of different major histocompatibility (MHC) backgrounds. To this end, the efficacy of such immunogens, in respect of eliciting an immune response against PV, might be expected
15 to differ between animals of diverse MHC background.

In addition, there is surprisingly little knowledge regarding which PV proteins are expressed by infected cells at various stages of differentiation, and hence it is not possible to predict which proteins will be responsible for defining appropriate immunological
20 targets.

The present invention provides a polyprotein construct comprising a plurality of PV early ORF proteins in one fused or linked construct to improve the efficacy of immune stimulation against PV infection and to avoid the need to define specific immunological
25 targets.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides as an isolated product, a polyprotein
30 construct comprising at least two amino acid sequences fused directly or indirectly

together, each of said sequences being the sequence of an early open reading frame (ORF) protein of papillomavirus (PV) or an immunogenic variant or fragment thereof, and at least one of said sequences being other than the E6 or E7 protein sequence or an immunogenic variant or fragment thereof.

5

In yet another aspect, the present invention provides a composition for eliciting a humoral and/or cellular immune response against PV in a host animal, said composition comprising an immunologically effective amount of a construct as described above, together with a pharmaceutically acceptable carrier and/or diluent.

10

In yet another aspect, this invention provides a method for eliciting a humoral and/or cellular response against PV in a host animal, which method comprises administering to the host animal an immunologically effective amount of a polyprotein construct as described above. In a related aspect, the invention also extends to use of
15 such a polyprotein construct in eliciting an immune response against PV in a host animal. Preferably, the host animal is a human, however the host animal may also be a non-human mammal.

The present invention also extends to a nucleic acid molecule which encodes a
20 polypeptide construct as broadly described above. Such a nucleic acid molecule may be delivered to a host animal in a nucleic acid vaccine composition with a pharmaceutically acceptable carrier and/or diluent, for expression of the encoded polyprotein construct *in vivo* in a host animal. Alternatively, the nucleic acid molecule may be included in a recombinant DNA molecule comprising an expression control sequence operatively
25 linked to the nucleic acid molecule.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of
30 integers but not the exclusion of any other integer or group of integers."

- 5 -

DETAILED DESCRIPTION OF THE INVENTION

The term "polyprotein construct" as used herein is used to describe a protein construct made up of individual proteins that have been joined together in a sequence
5 whereby they retain their original relevant biological activities.

The term "isolated" as used herein denotes that the polyprotein construct has undergone at least one purification or isolation step, and preferably is in a form suitable for administration to a host animal.

10

By use of the term "immunologically effective amount" herein in the context of treatment of PV infection, it is meant that the administration of that amount to an individual PV infected host, either in a single dose or as part of a series, that is effective for treatment of PV infection. By the use of the term "immunologically effective amount"
15 herein in the context of prevention of PV infection, it is meant that the administration of that amount to an individual host, either in a single dose or as part of a series, that is effective to delay, inhibit, treat or prevent PV infection or disease. The effective amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated, the capacity of the individual's immune
20 system to synthesise antibodies, the degree of protection desired, the formulation of the immunogen, the assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

25 Preferably, the amino acid sequences in the polyprotein construct substantially correspond to the sequences of wild-type early ORF proteins of PV, including allelic or other variants thereof. Suitable variants include variants having single or multiple amino acid substitutions or additions to the wild-type sequences, and may have at least 50-60%, more preferably at least 70-80%, and most preferably at least 90%, similarity to the wild-
30 type amino acid sequences, provided the variant is capable of eliciting an immune

response against PV in a host animal. The amino acid sequences may also be immunogenic fragments of the wild-type early ORF proteins, that is fragments of the proteins which are capable of eliciting an immune response in a host animal. Suitably, the immunogenic fragment will comprise at least five, and more preferably at least ten,
5 contiguous amino acid residues of the particular protein. Such immunogenic fragments may also be recognised by PV-specific antibodies, particularly antibodies which have a protective or therapeutic effect in relation to PV infection. Preferably, the immunogenic fragment is a non-full length fragment of a wild-type amino acid sequence, which may for example comprise a deletion mutant of an early ORF protein corresponding to at least
10 50%, more preferably 60-70%, and even 80-90% of the full length wild-type amino acid sequence.

The amino acid sequences in the polyprotein construct of the present invention may be selected from the group consisting of the E1, E2, E3, E4, E5 (E5a, E5b), E6, E7 and
15 E8 proteins of PV, and may be included in the construct in any desired order. By way of example, the construct may be selected from the group consisting of:

- (a) E6/E4
- (b) E6/E5a/E4
- (c) E6/E7/E4
- 20 (d) E6/E7/E5a/E4
- (e) E6/E7/E1/E4
- (f) E6/E7/E5a/E1/E4
- (g) E6/E7/E5a/E1/E2/E4
- (h) E6/E7/E5a/E5b/E1/E2/E4
- 25 (i) E2/E5b
- (j) E2/E1/E5b
- (k) E2/E5a/E5b
- (l) E2/E1/E5a/E5b
- (m) E2/E4/E5a/E5b/E6/E7/E1
- 30 (n) E2/E3/E4/E5/E8/E6/E7/E1.

- 7 -

As described above, at least one of the early ORF proteins is other than the E6 or E7 proteins. Preferably one of the early ORF proteins in the construct is the E4 protein.

The polyprotein constructs of this invention preferably comprise at least three, and
5 more preferably three, four or five early ORF protein sequences. In addition, two or more different polyprotein constructs based on different combinations of early ORF proteins and/or different PV genotypes may be included in a single composition for prophylactic or therapeutic use.

10 In the polyprotein constructs of this invention, the amino acid sequences may be fused or linked directly together. Alternatively, they may be linked with a linker sequence of from 1 to 50, preferably 1 to 20, and more preferably 1 to 5, amino acid residues between the separate amino acid sequences. By way of example, such a linker
15 comprising a restriction endonuclease site. Linker sequences as described above may also be provided before and/or after the amino acid sequences in the polyprotein constructs.

The polyprotein constructs of this invention may also comprise a tag protein or
20 peptide moiety fused or otherwise coupled thereto to assist in purification of the polyprotein construct. Suitable tag moieties include, for example, (His)₆, glutathione-S-transferase (GST) and FLAG (International Biotechnologies), with the (His)₆ tag moiety being preferred. The constructs may further comprise a component to enhance the immunogenicity of the polyprotein. The component may be an adjuvant such as
25 diphtheria or cholera toxin or *E. coli* heat labile toxin (LT), or a non-toxic derivative thereof such as the holotoxoid or B subunit of cholera toxin or LT. In addition, the polyprotein construct of the invention may comprise a lipid binding region to facilitate incorporation into ISCOMs. Suitable lipid binding regions are disclosed by way of example in Australian Provisional Patent Application No. PN8867/96, dated 25 March
30 1996. A preferred lipid binding region is an influenza haemagglutinin tail.

The present invention also provides a nucleic acid molecule comprising a sequence of nucleotides which encodes a polyprotein construct as broadly described above.

5 The nucleic acid molecule may be RNA or DNA, single stranded or double stranded, in linear or covalently closed circular form. It will be appreciated that the sequence of nucleotides of this aspect of the invention may be obtained from natural, synthetic or semi-synthetic sources; furthermore, this nucleotide sequence may be a naturally-occurring sequence, or it may be related by mutation, including single or
10 multiple base substitutions, deletions, insertions and inversions, to such a naturally-occurring sequence, provided always that the nucleic acid molecule comprising such a sequence is capable of being expressed as a polyprotein construct as described herein.

The nucleotide sequence may have expression control sequences positioned
15 adjacent to it, such control sequences being derived from either a homologous or a heterologous source.

Since nucleic acid molecules may be delivered directly as "naked DNA" to a host animal, (see, for example, Wolfe *et al.*, 1990, *Science* **247**:1465 and Fynan *et al.*, 1993,
20 *Proc.Natl. Acad. Sci. USA*, **90**:11478), the present invention also includes a nucleic acid vaccine composition comprising a nucleic acid molecule as described above, together with a pharmaceutically acceptable carrier and/or diluent.

Immunisation with an isolated nucleic acid molecule allows *in vivo* synthesis of
25 the encoded polyprotein construct by the host animal in a manner similar to the manner in which PV proteins are expressed during infection by PV. In this aspect, the present invention also extends to a method for eliciting an immune response against PV in a host animal, which method comprises administering to the host animal an immunologically effective amount of a nucleic acid molecule as described above. The invention also

extends to use of such a nucleic acid molecule in eliciting an immune response against PV in a host animal.

This invention also provides a recombinant DNA molecule comprising an
5 expression control sequence having promoter and initiator sequences, the nucleotide
sequence encoding the polyprotein construct being located 3' to the promoter and
initiator sequences and a terminator sequence located 3' to this sequence of nucleotides.
In yet another aspect, the invention provides a recombinant DNA cloning vehicle such
as a plasmid capable of expressing the polyprotein construct, as well as a host cell
10 containing a recombinant DNA cloning vehicle and/or a recombinant DNA molecule as
described above.

Suitable expression control sequences and host cell/cloning vehicle combinations
are well known in the art, and are described by way of example, in Sambrook *et al.*
15 (1989) *Molecular Cloning : A Laboratory Manual*, 2nd ed. Cold Spring Harbor, New
York, Cold Spring Harbor Laboratory Press. Thus, the nucleotide sequence may be
ligated into any suitable expression vector, which may be either a prokaryotic or
eukaryotic expression vector. Preferably, the vector is a prokaryotic expression vector
such as pTrcHisA or pGEX-STOP (a pGEX expression vector (Amrad/Pharmacia Biotech)
20 which has been manipulated so as to result in truncation of the GST moiety, disclosed in
Australian Provisional Patent Application No. PN8272/86, dated 26 February 1996).
Whilst the host cell is preferably a prokaryotic cell, more preferably a bacterium such as
E. coli, it will be understood that the host cell may alternatively be a yeast or other
eukaryotic cell, or insect cells infected with baculovirus or the like.

25

Once recombinant DNA cloning vehicles and/or host cells expressing a
polyprotein construct of this invention have been identified, the expressed polypeptides
synthesised by the host cells, for example, as a fusion protein, can be isolated
substantially free of contaminating host cell components by techniques well known to
30 those skilled in the art.

The polyprotein construct-encoding DNA sequence is formed by linking or "fusing" sequences encoding each of the individual protein moieties. The first sequence in the polyprotein DNA construction has a promoter element and a ribosome binding site. These elements assure that transcription of the polyprotein DNA into mRNA begins at a defined site and that the signal, the ribosome binding site, needed for translation of mRNA into protein is present. Synthesis of the polyprotein is made continuous from one protein component to the next by removing or altering any initiation or binding signals and stop codons from the subsequent protein-encoding sequences. The stop codon, normally a signal for the ribosome to stop translation and to end the polypeptide, is not altered or removed from the last DNA sequence. The individual protein encoding sequences are jointed such that a proper phasing is made of the mRNA reading frames for translation of the sequence into the desired amino acids. Once a DNA sequence encoding a polyprotein construct or a "polyprotein gene" is made, it is necessary to demonstrate that the construction leads to production of a stable polyprotein construct. If the resulting protein is not stable, for example because the junctions between the proteins are vulnerable to proteolytic digestion, then the junction regions are modified. This can be done by inserting different amino acids at or near the junction or by building spacers of amino acids between the individual proteins. Linkers or spacers can also be introduced to modify the overall activity of the polyprotein. By adjusting the space between and orientation of the individual proteins it is possible to modify the total activity of the polyprotein construct. Further details of the preparation of polyprotein constructs of the present invention by recombinant DNA techniques are disclosed, by way of example, in US Patent No. 4774180, the disclosure of which is incorporated herein by reference.

25

Preferably, the polymerase chain reaction (PCR) is used to amplify the nucleotide sequences encoding each of the individual PV early ORF proteins. The nucleotide sequences which are amplified may be full length or non full-length fragments thereof. Restriction endonuclease sites may be incorporated in the oligonucleotide primers used for PCR to furnish directional ligation of the amplification products in the same

30

- 11 -

translational frame and to enable directional cloning into a suitable expression vector. The primers may encode an artificial initiator codon or a termination codon.

The first nucleotide sequence has an initiator codon. This initiator codon may
5 either be the normal wild-type initiator codon of the first sequence or may be inserted artificially at another chosen position of this sequence. Synthesis of the polyprotein construct is made continuous from one protein component to the next by removing or altering any initiation or binding signals and termination codons. The termination codon must be present in the last nucleotide sequence. This is effected normally by not altering
10 or removing the termination codon of the last nucleotide sequence. However, this termination codon may be inserted artificially, by methods known to persons skilled in the art, by first removing the normal, wild-type termination codon of the last nucleotide sequence and inserting another, in the correct reading frame, at another position of this sequence.

15

The polyprotein construct-encoding DNA sequence may incorporate restriction sites at the flanking ends to facilitate insertion of the DNA sequence into a suitable expression vector.

20 The PV can be a human or an animal PV, and is preferably HPV. The HPV may be of any genotype, and may for example be selected from the group consisting of HPV-6, HPV-11, HPV-16, HPV-18, HPV-33, HPV-35, HPV-31 and HPV-45. Preferably, the HPV is HPV-6 or HPV-11.

25 The present invention is particularly, but not exclusively, directed to polyprotein constructs comprising early ORF proteins of the HPV-6 and HPV-11 genotypes which are causative agents of condylomata acuminata, however it will be appreciated that the invention extends to variants of the corresponding proteins in other HPV genotypes, particularly the HPV-16 and HPV-18 genotypes, and other genotypes which have
30 oncogenic potential of a type similar to HPV-16 and HPV-18.

- 12 -

The polyprotein constructs of the present invention may comprise early ORF proteins of a single HPV genotype, or alternatively they may comprise early ORF proteins from more than one HPV genotype. In addition, a combination of more than one polyprotein construct may be used in cases where not all early ORF proteins are represented in the one polyprotein construct, or where immune responses to more than one HPV genotype are desired.

The polyprotein constructs of the present invention are provided as isolated proteins, that is they are substantially free of other PV proteins, and find particular utility for the treatment of genital warts, cervical cancer or other conditions caused by HPV in man. The polyprotein constructs can be included in pharmaceutical compositions for the treatment or prevention of diseases involving HPV as well as the other conditions discussed above.

The polyprotein constructs of the invention may be used to raise antibodies and/or induce cellular immune responses, either in subjects for which protection against infection by PV is desired, i.e. as prophylactic vaccines, or to heighten the immune response to an PV infection already present, i.e. as therapeutic vaccines. They also can be injected into production species to obtain antisera. In lieu of the polyclonal antisera obtained in the production species, monoclonal antibodies may be produced using the standard methods or by more recent modifications thereof by immortalising spleen or other antibody-producing cells for injection into animals to obtain antibody-producing clones. The polyclonal or monoclonal antibodies obtained, corrected if necessary for species variations, can also be used as therapeutic agents.

Direct administration of the polyprotein constructs to a host animal such as a human can confer either protective immunity against PV or, if the subject is already infected, a boost to the subject's own immune response to more effectively combat the progress of the PV induced disease.

- 13 -

The magnitude of the prophylactic or therapeutic dose of a polyprotein constructs of this invention will, of course, vary with the group of patients (age, sex, etc.), the nature or the severity of the condition to be treated and with the particular polyprotein construct and its route of administration. In general, the weekly dose range for use lies within the
5 range of from about 0.1 to about 5 μ g per kg body weight of a mammal.

Any suitable route of administration may be employed for providing a mammal, especially a human, with an effective dosage of a polyprotein construct of this invention. For example, oral, rectal, vaginal, topical, parenteral, ocular, nasal, sublingual, buccal,
10 intravenous and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, suppositories, aerosols and the like. Said dosage forms also include injected or implanted slow releasing devices specifically designed for this purpose or other forms of implants modified to additionally act in this fashion.

15

If the polyprotein constructs are to be administered as vaccines, they are formulated according to conventional methods for such administration to the subject to be protected. The polyprotein constructs may be delivered in accordance with this invention in ISCOMSTM (immune stimulating complexes), liposomes or encapsulated in
20 compounds such as acrylates or poly(DL-lactide-co-glycoside) to form microspheres. They may also be incorporated into oily emulsions and delivered orally.

Other adjuvants, as well as conventional pharmaceutically acceptable carriers, excipients, buffers or diluents, may also be included in vaccine compositions of this
25 invention. Generally, a vaccine composition in accordance with the present invention will comprise an immunologically effective amount of the polyprotein construct, and optionally an adjuvant, in conjunction with one or more conventional pharmaceutically acceptable carriers and/or diluents. An extensive though not exhaustive list of adjuvants can be found in Coulter and Cox, "Advances in Adjuvant Technology and Application",
30 in *Animal Parasite Control Utilizing Biotechnology*, Chapter 4, Ed. Young, W.K., CRC

Press, 1992. As used herein "pharmaceutically acceptable carriers and/or diluents" include any and all solvents, dispersion media, aqueous solutions, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art and
5 is described by way of example in *Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Company, Pennsylvania, U.S.A.

In practical use, a polyprotein construct of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to
10 conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g. oral or parenteral (including intravenous and intra-arterial). In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water glycols, oils, alcohols, flavouring agents, preservatives, colouring agents
15 and the like in the case of oral liquid preparations, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations such as, for example, powders, capsules and tablets. Because of their ease of administration, tablets and capsules represent the most
20 advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar-coated or enteric-coated by standard techniques.

In addition to the common dosage forms set out above, the polyprotein constructs
25 of this invention may also be administered by controlled release means and/or delivery devices, including by way of example, the controlled release preparations disclosed in International Patent Specification No. PCT/AU93/00677 (Publication No. WO 94/15636).

Pharmaceutical compositions of the present invention suitable for oral or
30 parenteral administration may be presented as discrete units such as capsules, cachets or

- 15 -

tablets each containing a predetermined amount of the active ingredient, as a powder or granules or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion or a water-in-oil liquid emulsion. Such compositions may be prepared by any of the methods of pharmacy but all methods include the step of bringing
5 into association the active ingredient with the carrier which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation.

10 Further features of the present invention are more fully described in the following Example(s). It is to be understood, however, that this detailed description is included solely for the purposes of exemplifying the present invention, and should not be understood in any way as a restriction on the broad description of the invention as set out above.

15

EXAMPLES

Example 1 - Amplification and cloning of early open reading frames (ORFs) of HPV6b

20 A clone containing the entire genome of HPV6b in pBR322 (de Villiers, 1981, *J. Virol*, **40**:932) was used as the template for separate PCR amplifications of E6, E7, E5a, E5b, E1, E2 and E4 open reading frame (ORF) sequences.

Appropriate restriction enzyme recognition sequences were included in the
25 oligonucleotides used for amplification (Table I; 1-7) to allow sequential assembly of these amplified early gene sequences into a 'polyprotein' sequence as depicted in Figure 1A.

- 16 -

In this scheme, E6 was amplified with oligonucleotides containing a *Sma*I site at the 5' end and *Hind*III, *Nco*I and *Xba*I sites at the 3' end. As well, E4 was amplified with oligonucleotides containing *Xba*I, *Sac*I, *Kpn*I and *Spe*I sites 5' and a *Bgl*II site 3'.

5 These amplified fragments were cloned as *Sma*I/*Xba*I (E6) and *Xba*I/*Bgl*II (E4) (Figure 1B) in the vector pSP70 (Promega Corporation) which had been modified by the removal of an *Eco*RV/*Eco*RI fragment to contain a portion of the pGEM3Zf (Promega Corporation) polylinker - *Hind*III through *Eco*RI. As well, unwanted sites upstream of the *Sma*I site were removed by cleaving with *Sma*I/*Xho*I and insertion of a *Sma*I/*Sal*I/*Xho*I
10 linker to create the vector pSP70 (MOD).

The E6/E4 cassette was able to be removed by cleavage with *Sma*I/*Bgl*II and this was then cloned for expression into the pGEX-STOP vector which produces a non-fusion protein with a C-terminal six-histidine sequence for purification purposes.

15

Using the introduced restriction enzyme recognition sequences, other early ORF sequences were incorporated into the E6/E4 cassette cloned into pSP70 (MOD) and then the newly created cassette cloned as a *Sma*I/*Bgl*II fragment into pGEX-STOP.

20 In this manner polyprotein constructs containing E6/E5a/E4, E6/E7/E4, E6/E7/E5a/E4, E6/E7/E1/E4 and E6/E7/E5a/E1/E4 were assembled. Complete DNA sequence data for the first three constructs is included and sequence data across the junctions of E1 is included for the latter two. DNA sequencing revealed the *Spe*I site was inactivated by a single base change which occurred either during oligonucleotide
25 synthesis, PCR or cloning.

As well the tetrafusion construct of E6/E7/E5a/E4 was cloned for expression into pET23b (Novagen) by firstly subcloning the tetramer as a *Sma*I/*Bgl*II fragment into the *Sma*I/*Bam*HI sites of the vector pRIT2T (AMRAD Pharmacia Biotech). The tetramer was



- 17 -

then removed by restriction with *Sma*I and *Sa*II and cloned into the *Hinc*II/*Xho*I sites of the vector pET23b.

A further construct containing E2 and E5b, but which could also accommodate the
5 addition of E1 and E5a, was created by amplifying E2 with oligonucleotides containing a *Sma*I site at the 5' end and *Xba*I, *Nco*I, *Kpn*I and *Sac*I, sites at the 3' end (Table 1; 8) and with E5b amplified using oligonucleotides with an *Xba*I site 5' and *Xho*I, *Bgl*II sites 3' (Table 1; 9). These amplified fragments were then cloned into pSP70 (MOD) as depicted in Figure 1C.

10

Table 1

Oligonucleotides used for PCR		
Early gene	Forward	Reverse
1 E6	5'GCGCCCCGGGATGGAAAGTGC AAATGCCTC ^{3'} (SEQ ID No. 1)	5'GCGCTCTAGACCATGGAAGCT TGGGTAACATGTCTTCCATGC ^{3'} (SEQ ID. No.2)
2 E4	5'GCGCTCTAGAGAGCTCGGTACC ACTAGTGGAGCACCAACATTGG GAAG ^{3'} (SEQ ID No. 3)	5'GCGCAGATCTTAGGCGTAGCT GAACTGTTAC ^{3'} (SEQ ID No. 4)
3 E5a	5'GCGCCCATGGGAAGTGGTGCCT GTACAAATAGC ^{3'} (SEQ ID No. 5)	5'GCGCTCTAGATTGCTGTGTGG TAACAATATAG ^{3'} (SEQ ID No. 6)
4 E7	5'GCGCAAGCTTCATGGAAGACAT GTTACCCTAAAG ^{3'} (SEQ ID No. 7)	5'GCGCCCATGGGGTCTTCGGT GCGCAGATGG ^{3'} (SEQ ID No. 8)
5 E1	5'GCGCGAGCTCGCGGACGATTCA GGTACAGAAAATG ^{3'} (SEQ ID No. 9)	5'GCGCGGTACCTAAAGTTCTAA CAACTGTTCTG ^{3'} (SEQ ID No. 10)
6 E2	5'GCGCGGTACCGAAGCAATAGCC AAGCGTTTAG ^{3'} (SEQ ID No. 11)	5'GCGCACTAGTCAATAGGTGCA GTGACATAAATC ^{3'} (SEQ ID No. 12)
7 E5b	5'GCGCTCTAGACTAACATGTCAAT TTAATGATG ^{3'} (SEQ ID No. 13)	5'GCGCGAGCTCATTCATATATA TATAATCACC ^{3'} (SEQ ID No. 14)
8 E2	5'GCGCCCCGGGATGGAAGCAATA GCCAAGCG ^{3'} (SEQ ID No. 15)	5'GCGCTCTAGACCATGGGGTAC CGAGCTCCAATAGGTGCAGTG ACATAAATC ^{3'} (SEQ ID No. 16)
9 E5b	5'GCGCTCTAGACTAACATGTCAAT TTAATGATG ^{3'} (SEQ ID No. 17)	5'GCGCAGATCTCTCGAGATTCA TATATATATAATCAC ^{3'} (SEQ ID No. 18)



Example 2 - Expression of different polyprotein constructs

The following constructs in pGEX-STOP were expressed in *E. coli* strain BL21 and protein production was assayed by PAGE followed by Western blotting:

5

- i) E6/E4
- ii) E6/E5a/E4
- iii) E6/E7/E4
- iv) E6/E7/E5a/E4

10

Construct (iv) in pET23b, expressed in *E. coli* strains BL21(DE3)pLysS and AD494(DE3)pLysS (Novagen), was also assayed for protein production by Western blotting and also by Coomassie Blue staining for the latter strain.

15

Cultures of 200mL were grown in Terrific broth (Tartoff and Hobbs, *Focus*, 9: 12, 1987) in the presence of 100 μ g/mL ampicillin (BL21) and 34 μ g/ml chloramphenicol [BL21(DE3)pLysS] and 15 μ g/mL kanamycin [AD494(DE3)pLysS]. At OD₆₀₀ ~ 1 protein expression was induced by the addition of IPTG to 0.4mM. Following induction samples were taken at 1, 2, 3, 4 and 5 hours and in some cases after overnight culture.

20

Figure 2 shows a Western blot result for the E6/E4 construct. This was probed with a polyclonal rabbit anti-E4 antibody (MWE4 - raised to the peptide LGNEHEESNSPLATPCVWPT conjugated to ovalbumin). An immunoreactive band of ~ 30 kDa was present in the 4 hour-induced sample (lanes 2 & 4, arrow) which was not
25 present in the uninduced sample (lane 3).

The same ~ 30kDa band can also be seen in the induced sample in Figure 3, lane 3, arrow (lane 2-uninduced) while the E6/E5a/E4 trimer construct of ~ 40kDa was poorly represented after a 4 hour induction period (lane 5, arrow; uninduced sample-lane 4)
30 using the same anti-E4 antibody.

In contrast however, a trimer construct of E6/E7/E4 (~ 41 kDa) could be easily detected after 5 hours induction using an anti-hexahistidine monoclonal antibody (Dianova) [Figure 4, lane 4, arrow; uninduced sample - lane 3].

5 The same trimer construct was again easily visualised after 5 hours induction using the anti-E4 antibody MWE4 (Figure 5, lane TRI, arrow; control sample - lane C) and the tetramer consisting of E6/E7/E5a/E4 (~ 51 kDa) could also be detected (lane TET, arrow). Although this band is weak, it must be noted that a considerable amount of high molecular weight material is also immunoreactive, indicating the tetramer is reasonably
10 well expressed but possibly prone to aggregation.

Figure 6 indicates that an anti-E6 antibody (prepared as described below) was able to detect E6/E7/E4 after 5 hours induction (lane TRI, arrow) but not E6/E7/E5a/E4 (lane TET; lane C - uninduced). However, an anti-E7 antibody (prepared as described below)
15 was able to detect after 5 hours induction both the trimer (Figure 7, lane TRI, arrow; lane C - uninduced) and the tetramer (lane TET, arrow; lane C - uninduced), with the latter again showing indications of aggregation. A monoclonal antibody raised to an E4 peptide also recognised the trimer.

20 The phenomenon of aggregation was clearly apparent when the E6/E7/E5a/E4 tetramer was expressed in the pET23b plasmid in BL21(DE3)pLysS (Figure 8 - a Western blot probed with MWE4). Lanes 2-5 are 1 hour, 2 hour, 3 hour and overnight uninduced samples and lanes 6-9 represent 1 hour, 2 hour, 3 hour and overnight induced samples. After 1 hour induction a band of E6/E7/E5a/E4 can clearly be seen (arrow), but with
25 increased times of induction this seems to decrease and aggregated forms are increased. In contrast, when strain AD494(DE3)pLysS was used to express the tetramer, a substantial signal was obtained at the ~ 50kDa position on a Western blot of the insoluble fraction (Figure 9, arrow) following 2 hours induction, which still persisted at 3 hours. This immunoreactive band was not present in control samples and no protein was detected
30 in the samples from the soluble fractions.



- 21 -

Figure 10 shows the Coomassie stained profile of an identical gel, indicating that the immunoreactive bands present after 2 and 3 hours induction (Figure 9) can clearly be visualised as stained bands (arrow) which are not present in the control samples.

5 Example 3 - DNA sequencing of polyprotein constructs

Polyprotein constructs were sequenced in both directions by the dideoxy method using primers that generated overlapping sequence information. The ³²P-Sequencing™ Kit (Pharmacia) was used to generate ³⁵S-labelled chain-terminated fragments which were
10 analysed on a Sequi-Gen™ (Biorad) electrophoretic gel apparatus. The DNA and corresponding amino acid sequences for E6/E5a/E4 (CSL690.SEQ), E6/E7/E4 (CSL760.SEQ) and E6/E7/E5a/E4 (CSL673.SEQ) are shown below. (SEQ ID Nos: 19 and 20, 21 and 22, and 23 and 24, respectively).

15 For constructs E6/E7/E1/E4 (CSL 791) and E6/E7/E5a/E1/E4 (CSL 762), which were created from E6/E7/E4 and E6/E7/E5a/E4, respectively, DNA sequence analysis across the junctions of E1 with its neighbours is shown below (SEQ ID Nos. 25 and 26, 27 and 28, and 29 and 30, respectively).

- 22 -

File : CSL690.SEQ

Range : 1 - 11

Mode : Normal

Codon Table : Universal

E6/E5a/E4 - SEQ ID Nos, 19 (DNA) and 20 (amino acid)

	9	18	27	36	45	54												
5'	ATG	GAA	AGT	GCA	AAT	GCC	TCC	ACG	TCT	GCA	ACG	ACC	ATA	GAC	CAG	TTG	TGC	AAG
	Met	Glu	Ser	Ala	Asn	Ala	Ser	Thr	Ser	Ala	Thr	Thr	Ile	Asp	Gln	Leu	Cys	Lys
	63	72	81	90	99	108												
	ACG	TTT	AAT	CTA	TCT	ATG	CAT	ACG	TTG	CAA	ATT	AAT	TGT	GTG	TTT	TGC	AAG	AAT
	Thr	Phe	Asn	Leu	Ser	Met	His	Thr	Leu	Gln	Ile	Asn	Cys	Val	Phe	Cys	Lys	Asn
	117	126	135	144	153	162												
	GCA	CTG	ACC	ACA	GCA	GAG	ATT	TAT	TCA	TAT	GCA	TAT	AAA	CAC	CTA	AAG	GTC	CTG
	Ala	Leu	Thr	Thr	Ala	Glu	Ile	Tyr	Ser	Tyr	Ala	Tyr	Lys	His	Leu	Lys	Val	Leu
	171	180	189	198	207	216												
	TTT	CGA	GGC	GGC	TAT	CCA	TAT	GCA	GCC	TGC	GCG	TGC	TGC	CTA	GAA	TTT	CAT	GGA
	Phe	Arg	Gly	Gly	Tyr	Pro	Tyr	Ala	Ala	Cys	Ala	Cys	Cys	Leu	Glu	Phe	His	Gly
	225	234	243	252	261	270												
	AAA	ATA	AAC	CAA	TAT	AGA	CAC	TTT	GAT	TAT	GCT	GGA	TAT	GCA	ACA	ACA	GTT	GAA
	Lys	Ile	Asn	Gln	Tyr	Arg	His	Phe	Asp	Tyr	Ala	Gly	Tyr	Ala	Thr	Thr	Val	Glu
	279	288	297	306	315	324												
	GAA	GAA	ACT	AAA	CAA	GAC	ATC	TTA	GAC	GTG	CTA	ATT	CGG	TGC	TAC	CTG	TGT	CAC
	Glu	Glu	Thr	Lys	Gln	Asp	Ile	Leu	Asp	Val	Leu	Ile	Arg	Cys	Tyr	Leu	Cys	His
	333	342	351	360	369	378												
	AAA	CCG	CTG	TGT	GAA	GTA	GAA	AAG	GTA	AAA	CAT	ATA	CTA	ACC	AAG	GCG	CGG	TTC
	Lys	Pro	Leu	Cys	Glu	Val	Glu	Lys	Val	Lys	His	Ile	Leu	Thr	Lys	Ala	Arg	Phe
	387	396	405	414	423	432												
	ATA	AAG	CTA	AAT	TGT	ACG	TGG	AAG	GGT	CGC	TGC	CTA	CAC	TGC	TGG	ACA	ACA	TGC
	Ile	Lys	Leu	Asn	Cys	Thr	Trp	Lys	Gly	Arg	Cys	Leu	His	Cys	Trp	Thr	Thr	Cys
	441	450	459	468	477	486												
	ATG	GAA	GAC	ATG	TTA	CCC	AAG	CTT	CCA	TGG	GAA	GTG	GTG	CCT	GTA	CAA	ATA	GCT
	Met	Glu	Asp	Met	Leu	Pro	Lys	Leu	Pro	Trp	Glu	Val	Val	Pro	Val	Gln	Ile	Ala
	495	504	513	522	531	540												
	GCA	GGA	ACA	ACC	AGC	ACA	TTC	ATA	CTG	CCT	GTT	ATA	ATT	GCA	TTT	GTT	GTA	TGT
	Ala	Gly	Thr	Thr	Ser	Thr	Phe	Ile	Leu	Pro	Val	Ile	Ile	Ala	Phe	Val	Val	Cys
	549	558	567	576	585	594												

- 23 -

```

TTT GTT AGC ATC ATA CTT ATT GTA TGG ATA TCT GAG TTT ATT GTG TAC ACA TCT
---
Phe Val Ser Ile Ile Leu Ile Val Trp Ile Ser Glu Phe Ile Val Tyr Thr Ser

        603          612          621          630          639          648

GTG CTA GTA CTA ACA CTG CTT TTA TAT TTA CTA TTG TGG CTG CTA TTA ACA ACC
---
Val Leu Val Leu Thr Leu Leu Leu Tyr Leu Leu Leu Trp Leu Leu Leu Thr Thr

        657          666          675          684          693          702

CCC TTG CAA TTT TTC CTA CTA ACT CTA CTT GTG TGT TAC TGT CCC GCA TTG TAT
---
Pro Leu Gln Phe Phe Leu Leu Thr Leu Leu Val Cys Tyr Cys Pro Ala Leu Tyr

        711          720          729          738          747          756

ATA CAC TAC TAT ATT GTT ACC ACA CAG CAA TCT AGA GAG CTC GGT ACC ACT AAT
---
Ile His Tyr Tyr Ile Val Thr Thr Gln Gln Ser Arg Glu Leu Gly Thr Thr Asn

        765          774          783          792          801          810

GGA GCA CCA AAC ATT GGG AAG TAT GTT ATG GCA GCA CAG TTA TAT GTT CTC CTG
---
Gly Ala Pro Asn Ile Gly Lys Tyr Val Met Ala Ala Gln Leu Tyr Val Leu Leu

        819          828          837          846          855          864

CAT CTG TAT CTA GCA CTA CAC AAG AAG TAT CCA TTC CTG AAT CTA CTA CAT ACA
---
His Leu Tyr Leu Ala Leu His Lys Lys Tyr Pro Phe Leu Asn Leu Leu His Thr

        873          882          891          900          909          918

CCC CCG CAC AGA CCT CCA CCC TTG TGT CCT CAA GCA CCA AGG AAG ACG CAG TGC
---
Pro Pro His Arg Pro Pro Pro Leu Cys Pro Gln Ala Pro Arg Lys Thr Gln Cys

        927          936          945          954          963          972

AAA CGC CGC CTA GGA AAC GAG CAC GAG GAG TCC AAC AGT CCC CTT GCA ACG CCT
---
Lys Arg Arg Leu Gly Asn Glu His Glu Glu Ser Asn Ser Pro Leu Ala Thr Pro

        981          990          999          1008          1017          1026

TGT GTG TGG CCC ACA TTG GAC CCG TGG ACA GTG GAA ACC ACA ACC TCA TCA CTA
---
Cys Val Trp Pro Thr Leu Asp Pro Trp Thr Val Glu Thr Thr Thr Ser Ser Leu

        1035          1044          1053          1062          1071          1080

ACA ATC ACG ACC AGC ACC AAA GAC GGA ACA ACA GTA ACA GTT CAG CTA CGC CTA
---
Thr Ile Thr Thr Ser Thr Lys Asp Gly Thr Thr Val Thr Val Gln Leu Arg Leu

        1089          1098          1107

AGA TCT CAT CAC CAT CAC CAT CAC TAA 3'
---
Arg Ser His His His His His His ***

```

- 24 -

File : CSL760.SEQ

Range : 1 - 1128 Mode : Normal

Codon Table : Universal

E6/E7/E4 - SEQ ID Nos. 21 (DNA) and 22 (amino acid)

	9	18	27	36	45	54												
5'	ATG	GAA	AGT	GCA	AAT	GCC	TCC	ACG	TCT	GCA	ACG	ACC	ATA	GAC	CAG	TTG	TGC	AAG
	Met	Glu	Ser	Ala	Asn	Ala	Ser	Thr	Ser	Ala	Thr	Thr	Ile	Asp	Gln	Leu	Cys	Lys
	63	72	81	90	99	108												
	ACG	TTT	AAT	CTA	TCT	ATG	CAT	ACG	TTG	CAA	ATT	AAT	TGT	GTG	TTT	TGC	AAG	AAT
	Thr	Phe	Asn	Leu	Ser	Met	His	Thr	Leu	Gln	Ile	Asn	Cys	Val	Phe	Cys	Lys	Asn
	117	126	135	144	153	162												
	GCA	CTG	ACC	ACA	GCA	GAG	ATT	TAT	TCA	TAT	GCA	TAT	AAA	CAC	CTA	AAG	GTC	CTG
	Ala	Leu	Thr	Thr	Ala	Glu	Ile	Tyr	Ser	Tyr	Ala	Tyr	Lys	His	Leu	Lys	Val	Leu
	171	180	189	198	207	216												
	TTT	CGA	GGC	GGC	TAT	CCA	TAT	GCA	GCC	TGC	GCG	TGC	TGC	CTA	GAA	TTT	CAT	GGA
	Phe	Arg	Gly	Gly	Tyr	Pro	Tyr	Ala	Ala	Cys	Ala	Cys	Cys	Leu	Glu	Phe	His	Gly
	225	234	243	252	261	270												
	AAA	ATA	AAC	CAA	TAT	AGA	CAC	TTT	GAT	TAT	GCT	GGA	TAT	GCA	ACA	ACA	GTT	GAA
	Lys	Ile	Asn	Gln	Tyr	Arg	His	Phe	Asp	Tyr	Ala	Gly	Tyr	Ala	Thr	Thr	Val	Glu
	279	288	297	306	315	324												
	GAA	GAA	ACT	AAA	CAA	GAC	ATC	TTA	GAC	GTG	CTA	ATT	CGG	TGC	TAC	CTG	TGT	CAC
	Glu	Glu	Thr	Lys	Gln	Asp	Ile	Leu	Asp	Val	Leu	Ile	Arg	Cys	Tyr	Leu	Cys	His
	333	342	351	360	369	378												
	AAA	CCG	CTG	TGT	GAA	GTA	GAA	AAG	GTA	AAA	CAT	ATA	CTA	ACC	AAG	GCG	CGG	TTC
	Lys	Pro	Leu	Cys	Glu	Val	Glu	Lys	Val	Lys	His	Ile	Leu	Thr	Lys	Ala	Arg	Phe
	387	396	405	414	423	432												
	ATA	AAG	CTA	AAT	TGT	ACG	TGG	AAG	GGT	CGC	TGC	CTA	CAC	TGC	TGG	ACA	ACA	TGC
	Ile	Lys	Leu	Asn	Cys	Thr	Trp	Lys	Gly	Arg	Cys	Leu	His	Cys	Trp	Thr	Thr	Cys
	441	450	459	468	477	486												
	ATG	GAA	GAC	ATG	TTA	CCC	AAG	CTT	CAT	GGA	AGA	CAT	GTT	ACC	CTA	AAG	GAT	ATT
	Met	Glu	Asp	Met	Leu	Pro	Lys	Leu	His	Gly	Arg	His	Val	Thr	Leu	Lys	Asp	Ile
	495	504	513	522	531	540												
	GTA	TTA	GAC	CTG	CAA	CCT	CCA	GAC	CCT	GTA	GGG	TTA	CAT	TGC	TAT	GAG	CAA	TTA
	Val	Leu	Asp	Leu	Gln	Pro	Pro	Asp	Pro	Val	Gly	Leu	His	Cys	Tyr	Glu	Gln	Leu
	549	558	567	576	585	594												

- 25 -

GTA	GAC	AGC	TCA	GAA	GAT	GAG	GTG	GAC	GAA	GTG	GAC	GGA	CAA	GAT	TCA	CAA	CCT
Val	Asp	Ser	Ser	Glu	Asp	Glu	Val	Asp	Glu	Val	Asp	Gly	Gln	Asp	Ser	Gln	Pro
603				612				621				630			639		648
TTA	AAA	CAA	CAT	TTC	CAA	ATA	GTG	ACC	TGT	TGC	TGT	GGA	TGT	GAC	AGC	AAC	GTT
Leu	Lys	Gln	His	Phe	Gln	Ile	Val	Thr	Cys	Cys	Cys	Gly	Cys	Asp	Ser	Asn	Val
657				666				675				684			693		702
CGA	CTG	GTT	GTG	CAG	TGT	ACA	GAA	ACA	GAC	ATC	AGA	GAA	GTG	CAA	CAG	CTT	CTG
Arg	Leu	Val	Val	Gln	Cys	Thr	Glu	Thr	Asp	Ile	Arg	Glu	Val	Gln	Gln	Leu	Leu
711				720				729				738			747		756
TTG	GGA	ACA	CTA	AAC	ATA	GTG	TGT	CCC	ATC	TGC	GCA	CCG	AAG	ACC	CCA	TGG	TCT
Leu	Gly	Thr	Leu	Asn	Ile	Val	Cys	Pro	Ile	Cys	Ala	Pro	Lys	Thr	Pro	Trp	Ser
765				774				783				792			801		810
AGA	GAG	CTC	GGT	ACC	ACT	AAT	GGA	GCA	CCA	AAC	ATT	GGG	AAG	TAT	GTT	ATG	GCA
Arg	Glu	Leu	Gly	Thr	Thr	Asn	Gly	Ala	Pro	Asn	Ile	Gly	Lys	Tyr	Val	Met	Ala
819				828				837				846			855		864
GCA	CAG	TTA	TAT	GTT	CTC	CTG	CAT	CTG	TAT	CTA	GCA	CTA	CAC	AAG	AAG	TAT	CCA
Ala	Gln	Leu	Tyr	Val	Leu	Leu	His	Leu	Tyr	Leu	Ala	Leu	His	Lys	Lys	Tyr	Pro
873				882				891				900			909		918
TTC	CTG	AAT	CTA	CTA	CAT	ACA	CCC	CCG	CAC	AGA	CCT	CCA	CCC	TTG	TGT	CCT	CAA
Phe	Leu	Asn	Leu	Leu	His	Thr	Pro	Pro	His	Arg	Pro	Pro	Pro	Leu	Cys	Pro	Gln
927				936				945				954			963		972
GCA	CCA	AGG	AAG	ACG	CAG	TGC	AAA	CGC	CGC	CTA	GGA	AAC	GAG	CAC	GAG	GAG	TCC
Ala	Pro	Arg	Lys	Thr	Gln	Cys	Lys	Arg	Arg	Leu	Gly	Asn	Glu	His	Glu	Glu	Ser
981				990				999				1008			1017		1026
AAC	AGT	CCC	CTT	GCA	ACG	CCT	TGT	GTG	TGG	CCC	ACA	TTG	GAC	CCG	TGG	ACA	GTG
Asn	Ser	Pro	Leu	Ala	Thr	Pro	Cys	Val	Trp	Pro	Thr	Leu	Asp	Pro	Trp	Thr	Val
1035				1044				1053				1062			1071		1080
GAA	ACC	ACA	ACC	TCA	TCA	CTA	ACA	ATC	ACG	ACC	AGC	ACC	AAA	GAC	GGA	ACA	ACA
Glu	Thr	Thr	Thr	Ser	Ser	Leu	Thr	Ile	Thr	Thr	Ser	Thr	Lys	Asp	Gly	Thr	Thr
1089				1098				1107				1116			1125		
GTA	ACA	GTT	CAG	CTA	CGC	CTA	AGA	TCT	CAT	CAC	CAT	CAC	CAT	CAC	TAA	3'	
Val	Thr	Val	Gln	Leu	Arg	Leu	Arg	Ser	His	His	His	His	His	His	His	***	

- 26 -

File : CSL673.DNA

Range : 1 - 1377 Mode : Normal

Codon Table : Universal

E6/E7/E5a/E4 - SEQ ID Nos. 23 (DNA) and 24 (amino acid)

	9	18	27	36	45	54
5'	ATG GAA AGT GCA AAT GCC TCC ACG TCT GCA ACG ACC ATA GAC CAG TTG TGC AAG					
	Met Glu Ser Ala Asn Ala Ser Thr Ser Ala Thr Thr Ile Asp Gln Leu Cys Lys					
	63	72	81	90	99	108
	ACG TTT AAT CTA TCT ATG CAT ACG TTG CAA ATT AAT TGT GTG TTT TGC AAG AAT					
	Thr Phe Asn Leu Ser Met His Thr Leu Gln Ile Asn Cys Val Phe Cys Lys Asn					
	117	126	135	144	153	162
	GCA CTG ACC ACA GCA GAG ATT TAT TCA TAT GCA TAT AAA CAC CTA AAG GTC CTG					
	Ala Leu Thr Thr Ala Glu Ile Tyr Ser Tyr Ala Tyr Lys His Leu Lys Val Leu					
	171	180	189	198	207	216
	TTT CGA GGC GGC TAT CCA TAT GCA GCC TGC GCG TGC TGC CTA GAA TTT CAT GGA					
	Phe Arg Gly Gly Tyr Pro Tyr Ala Ala Cys Ala Cys Cys Leu Glu Phe His Gly					
	225	234	243	252	261	270
	AAA ATA AAC CAA TAT AGA CAC TTT GAT TAT GCT GGA TAT GCA ACA ACA GTT GAA					
	Lys Ile Asn Gln Tyr Arg His Phe Asp Tyr Ala Gly Tyr Ala Thr Thr Val Glu					
	279	288	297	306	315	324
	GAA GAA ACT AAA CAA GAC ATC TTA GAC GTG CTA ATT CGG TGC TAC CTG TGT CAC					
	Glu Glu Thr Lys Gln Asp Ile Leu Asp Val Leu Ile Arg Cys Tyr Leu Cys His					
	333	342	351	360	369	378
	AAA CCG CTG TGT GAA GTA GAA AAG GTA AAA CAT ATA CTA ACC AAG GCG CGG TTC					
	Lys Pro Leu Cys Glu Val Glu Lys Val Lys His Ile Leu Thr Lys Ala Arg Phe					
	387	396	405	414	423	432
	ATA AAG CTA AAT TGT ACG TGG AAG GGT CGC TGC CTA CAC TGC TGG ACA ACA TGC					
	Ile Lys Leu Asn Cys Thr Trp Lys Gly Arg Cys Leu His Cys Trp Thr Thr Cys					
	441	450	459	468	477	486
	ATG GAA GAC ATG TTA CCC AAG CTT CAT GGA AGA CAT GTT ACC CTA AAG GAT ATT					
	Met Glu Asp Met Leu Pro Lys Leu His Gly Arg His Val Thr Leu Lys Asp Ile					
	495	504	513	522	531	540
	GTA TTA GAC CTG CAA CCT CCA GAC CCT GTA GGG TTA CAT TGC TAT GAG CAA TTA					
	Val Leu Asp Leu Gln Pro Pro Asp Pro Val Gly Leu His Cys Tyr Glu Gln Leu					
	549	558	567	576	585	594

- 27 -

GTA	GAC	AGC	TCA	GAA	GAT	GAG	GTG	GAC	GAA	GTG	GAC	GGA	CAA	GAT	TCA	CAA	CCT
Val	Asp	Ser	Ser	Glu	Asp	Glu	Val	Asp	Glu	Val	Asp	Gly	Gln	Asp	Ser	Gln	Pro
		603			612			621			630			639			648
TTA	AAA	CAA	CAT	TTC	CAA	ATA	GTG	ACC	TGT	TGC	TGT	GGA	TGT	GAC	AGC	AAC	GTT
Leu	Lys	Gln	His	Phe	Gln	Ile	Val	Thr	Cys	Cys	Cys	Gly	Cys	Asp	Ser	Asn	Val
		657			666			675			684			693			702
CGA	CTG	GTT	GTG	CAG	TGT	ACA	GAA	ACA	GAC	ATC	AGA	GAA	GTG	CAA	CAG	CTT	CTG
Arg	Leu	Val	Val	Gln	Cys	Thr	Glu	Thr	Asp	Ile	Arg	Glu	Val	Gln	Gln	Leu	Leu
		711			720			729			738			747			756
TTG	GGA	ACA	CTA	AAC	ATA	GTG	TGT	CCC	ATC	TGC	GCA	CCG	AAG	ACC	CCA	TGG	GAA
Leu	Gly	Thr	Leu	Asn	Ile	Val	Cys	Pro	Ile	Cys	Ala	Pro	Lys	Thr	Pro	Trp	Glu
		765			774			783			792			801			810
GTG	GTG	CCT	GTA	CAA	ATA	GCT	GCA	GGA	ACA	ACC	AGC	ACA	TTC	ATA	CTG	CCT	GTT
Val	Val	Pro	Val	Gln	Ile	Ala	Ala	Gly	Thr	Thr	Ser	Thr	Phe	Ile	Leu	Pro	Val
		819			828			837			846			855			864
ATA	ATT	GCA	TTT	GTT	GTA	TGT	TTT	GTT	AGC	ATC	ATA	CTT	ATT	GTA	TGG	ATA	TCT
Ile	Ile	Ala	Phe	Val	Val	Cys	Phe	Val	Ser	Ile	Ile	Leu	Ile	Val	Trp	Ile	Ser
		873			882			891			900			909			918
GAG	TTT	ATT	GTG	TAC	ACA	TCT	GTG	CTA	GTA	CTA	ACA	CTG	CTT	TTA	TAT	TTA	CTA
Glu	Phe	Ile	Val	Tyr	Thr	Ser	Val	Leu	Val	Leu	Thr	Leu	Leu	Leu	Tyr	Leu	Leu
		927			936			945			954			963			972
TTG	TGG	CTG	CTA	TTA	ACA	ACC	CCC	TTG	CAA	TTT	TTC	CTA	CTA	ACT	CTA	CTT	GTG
Leu	Trp	Leu	Leu	Leu	Thr	Thr	Pro	Leu	Gln	Phe	Phe	Leu	Leu	Thr	Leu	Leu	Val
		981			990			999			1008			1017			1026
TGT	TAC	TGT	CCC	GCA	TTG	TAT	ATA	CAC	TAC	TAT	ATT	GTT	ACC	ACA	CAG	CAA	TCT
Cys	Tyr	Cys	Pro	Ala	Leu	Tyr	Ile	His	Tyr	Tyr	Ile	Val	Thr	Thr	Gln	Gln	Ser
		1035			1044			1053			1062			1071			1080
AGA	GAG	CTC	GGT	ACC	ACT	AAT	GGA	GCA	CCA	AAC	ATT	GGG	AAG	TAT	GTT	ATG	GCA
Arg	Glu	Leu	Gly	Thr	Thr	Asn	Gly	Ala	Pro	Asn	Ile	Gly	Lys	Tyr	Val	Met	Ala
		1089			1098			1107			1116			1125			1134
GCA	CAG	TTA	TAT	GTT	CTC	CTG	CAT	CTG	TAT	CTA	GCA	CTA	CAC	AAG	AAG	TAT	CCA
Ala	Gln	Leu	Tyr	Val	Leu	Leu	His	Leu	Tyr	Leu	Ala	Leu	His	Lys	Lys	Tyr	Pro
		1143			1152			1161			1170			1179			1188

- 28 -

```

TTC CTG AAT CTA CTA CAT ACA CCC CCG CAC AGA CCT CCA CCC TTG TGT CCT CAA
---
Phe Leu Asn Leu Leu His Thr Pro Pro His Arg Pro Pro Pro Leu Cys Pro Gln
      1197          1206          1215          1224          1233          1242

GCA CCA AGG AAG ACG CAG TGC AAA CGC CGC CTA GGA AAC GAG CAC GAG GAG TCC
---
Ala Pro Arg Lys Thr Gln Cys Lys Arg Arg Leu Gly Asn Glu His Glu Glu Ser
      1251          1260          1269          1278          1287          1296

AAC AGT CCC CTT GCA ACG CCT TGT GTG TGG CCC ACA TTG GAC CCG TGG ACA GTG
---
Asn Ser Pro Leu Ala Thr Pro Cys Val Trp Pro Thr Leu Asp Pro Trp Thr Val
      1305          1314          1323          1332          1341          1350

GAA ACC ACA ACC TCA TCA CTA ACA ATC ACG ACC AGC ACC AAA GAC GGA ACA ACA
---
Glu Thr Thr Thr Ser Ser Leu Thr Ile Thr Thr Ser Thr Lys Asp Gly Thr Thr
      1359          1368          1377          1386          1395

GTA ACA GTT CAG CTA CGC CTA AGA TCT CAT CAC CAT CAC CAT CAC TAA 3'
---
Val Thr Val Gln Leu Arg Leu Arg Ser His His His His His His His ***

```

Junction of E1 and E4 ORFs for CSL791 and CSL762

SEQ ID Nos. 25 (DNA) and 26 (amino acid)

Modified
 Apn1 Sma1

5' GAG GAA GAT GGA AGC AAT AGC CAA GCG TTT AGA TGC GTG CCA GGA ACA GTT GTT AGA ACT TTA GGT ACC ACT AAT GGA GCA CCA AAC ATT GGG AAG TAT GTT ATG GCA 3'
 Glu Glu Asp Gly Ser Asn Ser Gln Ala Phe Arg Cys Val Pro Gly Thr Val Val Arg Thr Thr Asn Gly Ala Pro Asn Ile Gly Lys Tyr Val Met Ala

E1

E4

Junction of E5a and E1 for CSL762

SEQ ID Nos. 27 (DNA) and 28 (amino acid)

Xba1 Sac1

5' TGT CCC GCA TTG TAT ATA CAC TAC TAT ATT GTT ACC ACA CAG CAA TCT AGA GAG CTC GCG GAC GAT TCA GGT ACA GAA AAT GAG GGG TCT GGG TGT ACA GGA 3'
 Cys Pro Ala Leu Tyr Ile His Tyr Tyr Ile Val Thr Thr Gln Gln Ser Arg Glu Leu Ala Asp Asp Ser Gly Thr Glu Asn Glu Gly Ser Gly Cys Thr Gly

E5a

E1

Junction of E7 and E1 for CSL791

SEQ ID Nos. 29 (DNA) and 30 (amino acid)

Nco1 Xba1 Sac1

5' TTG GGA ACA CTA AAC ATA GTG TGT CCC ATC TGC GCA CCG AAG ACC CCA TGG TCT AGA GAG CTC GCG GAC GAT TCA GGT ACA GAA AAT GAG GGG TCT GGG TGT ACA GGA 3'
 Leu Gly Thr Leu Asn Ile Val Cys Pro Ile Cys Ala Pro Lys Thr Pro Trp Ser Arg Glu Leu Ala Asp Asp Ser Gly Thr Glu Asn Glu Gly Ser Gly Cys Thr Gly

E7

E1

- 30 -

Example 4 - Preparation of antibodies to HPV6b early ORF protein products

The following peptides corresponding to portions of the sequence of the relevant E proteins, were synthesised and conjugated to diphtheria toxoid:

5

E6 dip. tox-C-QYRHFDYAQYATTVEEETKQDILD
E7 MHGRHVTLKDIVLDLQPPD-C-dip. tox

For the E6 peptide two rabbits (following pre-bleeding) were each inoculated with approximately 54 μ g peptide/104 μ g diphtheria toxoid in Freund's complete adjuvant followed at 3-weekly intervals by a similar dose of peptide conjugate in Freund's incomplete adjuvant. Bleeds were taken one week after the second dose and one week following the third dose. The same regime was used for the E7 peptide using 45 μ g peptide/103 μ g diphtheria toxoid.

15

Serum derived from the bleeds were tested for specific antibody in a solid phase EIA against biotin-conjugated peptide which had been bound to plates coated with strepavidin.

20 Example 5 - Purification of polyprotein E6/E7/E4

The trimer polyprotein E6/E7/E4 was expressed in *E. coli* BL21 cells by induction of cells at OD₆₀₀ ~ 1 using 0.4mM IPTG. The cells were harvested by centrifugation (4,000g, 20 minutes), resuspended in 30mM Tris pH8.0, disrupted by sonication (MSE, amplitude 18 μ m, 4 x 30 seconds) and inclusion bodies pelleted by centrifugation (12,000g, 30 minutes). The pellet containing the trimer was solubilized in 8M Urea, 30mM Tris pH8.0 for 16 hours in the presence of protease inhibitors (Boehringer Cat. No. 1697498) and then centrifuged at 12,000g for 30 minutes and the supernatant collected. To this, Tris-(2-carboxyethyl)phosphine (TCEP) [Pierce] was added to 1.2mM final concentration. The supernatant was applied to Q-sepharose HP (Pharmacia) and the

- 31 -

column washed with one column volume of 8M Urea, 1.2mM TCEP, 30 mM Tris pH8.0. Fractions were then eluted using a gradient containing 0 to 1M NaCl in 10 column volumes of the washing buffer. The fractions obtained were examined by Western blot from 4 to 20% SDS-PAGE probed with the anti-E4 antibody MWE4.

5

Figure 11 shows a Western blot of material obtained from Q-sepharose. An immunoreactive band of ~ 41kDa is evident in supernatant material from the urea solubilisation lane 3, and in fractions corresponding to 120 to 150 mM NaCl (lanes 8 and 9, arrow).

10

Supernatant from the urea solubilisation was also applied to a column containing Chelating Sepharose Fast Flow (Pharmacia) to take advantage of the C-terminal six histidine sequence. Relatively poor binding of the trimer to the Nickel column was observed under the conditions described. The trimer was eluted from the column using a 0 to 500 mM imidazole gradient.

Example 6

In a further example of the present invention, a DNA sequence coding for a single polyprotein (Fig. 12) is formed by fusion of DNA fragments encoding HPV-6 early ORF proteins wherein the order of the ORFs is E2, E4, E5a, E5b, E6, E7 and E1.

The DNA sequences encoding the early ORF proteins are amplified individually by PCR using HPV-6 genomic DNA using the primers set out in Table 2.

25

Table 2

Gene	Oligonucleotides	
E2	(a)	5'-GTG TGT GAG CTC ATG GAA GCA ATA GCC AAG-3' (SEQ ID No. 31) and
	(b)	5'-GTG TGT GTC GAC CAA TAG GTG CAG TGA CAT-3' (SEQ ID No. 32)
E4	(c)	5'-GTG TGT GTC GAC ATG GGA GCA CCA AAC ATT-3' (SEQ ID No. 33) and
		5'-GTG TGT AGA TCT TAG GCG TAG CTG AAC TGT-3' (SEQ ID No. 34)
E5a	(e)	5'-GTG TGT AGA TCT ATG GAA GTG GTG CCT GTA-3' (SEQ ID No. 35) and
	(f)	5'-GTG TGT CTT AAG TTG CTG TGT GGT AAC AAT-3' (SEQ ID No. 36)
E5b	(g)	5'-GTG TGT CTT AAG ATG ATG CTA ACA TGT CAA-3' (SEQ ID No. 37) and
	(h)	5'-GTG TGT CCG CGG ATT CAT ATA TAT ATA ATC-3' (SEQ ID No. 38)
E6	(i)	5'-GTG TGT CCG CGG ATG GAA AGT GCA AAT GCC-3' (SEQ ID No. 39) and
	(j)	5'-GTG TGT GCT AGC GGG TAA CAT GTC TTC CTA-3' (SEQ ID No. 40)
E7	(k)	5'-GTG TGT GCT AGC ATG CAT GGA AGA CAT GTT-3' (SEQ ID No. 41) and
	(l)	5'-GTG TGT CGA TCG GGT CTT CGG TGC GCA GAT-3' (SEQ ID No. 42)
E1	(m)	5'-GTG TGT CGA TCG ATG GCG GAC GAT TCA GGT-3' (SEQ ID No. 43) and
	(n)	5'-GTG TGT GGT ACC TCA TAA AGT TCT AAC AAC-3' (SEQ ID No. 44)

The primers are synthesised to incorporate artificial restriction enzyme sites at the 5' and 3' termini of the amplification products. These restriction enzyme sites are used to facilitate the fusion of PCR products encoding the appropriate early ORF proteins in the desired order and in the correct translational frame. The restriction enzyme sites are also used to aid the cloning of the PCR products into the expression vector pTrcHisA. When cloned into this vector, the polyprotein construct is expressed as an N-terminal

- 33 -

hexaHis fusion. The nucleotide sequence and deduced amino acid sequence of this fusion are shown below (SEQ ID Nos. 45 and 46, respectively).

INFORMATION FOR HEXAHIS-POLYPROTEIN FUSION SEQUENCE:

- (I) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4770 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: HUMAN PAPILLOMAVIRUS TYPE 6
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION:1..4761
 - (D) OTHER INFORMATION:/codon_start= 1
/product= "HPV-6 Polyprotein"
- (ix) FEATURE:
 - (A) NAME/KEY: misc RNA
 - (B) LOCATION:1..108
 - (D) OTHER INFORMATION:/function= "Tag used for protein purification"
/product= "hexaHis leader sequence from pTrcHisA"
- (ix) FEATURE:
 - (A) NAME/KEY: misc_feature
 - (B) LOCATION:109..114
 - (D) OTHER INFORMATION:/label= SacI
- (ix) FEATURE:
 - (A) NAME/KEY: mRNA
 - (B) LOCATION:115..1218
 - (D) OTHER INFORMATION:/gene= "HPV-6 E2"
- (ix) FEATURE:
 - (A) NAME/KEY: misc_feature
 - (B) LOCATION:1219..1224
 - (D) OTHER INFORMATION:/label= SalI
- (ix) FEATURE:
 - (A) NAME/KEY: mRNA
 - (B) LOCATION:1225..1551
 - (D) OTHER INFORMATION:/gene= "HPV-6 E4"
- (ix) FEATURE:
 - (A) NAME/KEY: misc_feature
 - (B) LOCATION:1552..1557
 - (D) OTHER INFORMATION:/label= BglII
- (ix) FEATURE:
 - (A) NAME/KEY: mRNA
 - (B) LOCATION:1558..1830
 - (D) OTHER INFORMATION:/gene= "HPV-6 E5a"
- (ix) FEATURE:
 - (A) NAME/KEY: misc_feature
 - (B) LOCATION:1831..1836
 - (D) OTHER INFORMATION:/label= BfrI
- (ix) FEATURE:
 - (A) NAME/KEY: mRNA
 - (B) LOCATION:1837..2052
 - (D) OTHER INFORMATION:/gene= "HPV-6 E5b"

- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION:2053..2058
 (D) OTHER INFORMATION:/label= SacII
- (ix) FEATURE:
 (A) NAME/KEY: mRNA
 (B) LOCATION:2059..2508
 (D) OTHER INFORMATION:/gene= "HPV-6 E6"
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION:2509..2514
 (D) OTHER INFORMATION:/label= NheI
- (ix) FEATURE:
 (A) NAME/KEY: mRNA
 (B) LOCATION:2515..2808
 (D) OTHER INFORMATION:/gene= "HPV-6 E7"
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION:2809..2814
 (D) OTHER INFORMATION:/label= PvuI
- (ix) FEATURE:
 (A) NAME/KEY: mRNA
 (B) LOCATION:2815..4764
 (D) OTHER INFORMATION:/gene= "HPV-6 E1"
- (ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION:4765..4770
 (D) OTHER INFORMATION:/label= KpnI

ATG GGG GGT TCT CAT CAT CAT CAT CAT GGT ATG GCT AGC ATG ACT	48
Met Gly Gly Ser His His His His His Gly Met Ala Ser Met Thr	
1 5 10 15	
GGT GGA CAG CAA ATG GGT CGG GAT CTG TAC GAC GAT GAC GAT AAG GAT	96
Gly Gly Gln Gln Met Gly Arg Asp Leu Tyr Asp Asp Asp Asp Lys Asp	
20 25 30	
CGA TGG GGA TCC GAG CTC ATG GAA GCA ATA GCC AAG CGT TTA GAT GCG	144
Arg Trp Gly Ser Glu Leu Met Glu Ala Ile Ala Lys Arg Leu Asp Ala	
35 40 45	
TGC CAG GAA CAG TTG TTA GAA CTT TAT GAA GAA AAC AGT ACT GAC CTA	192
Cys Gln Glu Gln Leu Leu Glu Leu Tyr Glu Glu Asn Ser Thr Asp Leu	
50 55 60	
CAC AAA CAT GTA TTG CAT TGG AAA TGC ATG AGA CAT GAA AGT GTA TTA	240
His Lys His Val Leu His Trp Lys Cys Met Arg His Glu Ser Val Leu	
65 70 75 80	
TTA TAT AAA GCA AAA CAA ATG GGC CTA AGC CAC ATA GGA ATG CAA GTA	288
Leu Tyr Lys Ala Lys Gln Met Gly Leu Ser His Ile Gly Met Gln Val	
85 90 95	
GTG CCA CCA TTA AAG GTG TCC GAA GCA AAA GGA CAT AAT GCC ATT GAA	336
Val Pro Pro Leu Lys Val Ser Glu Ala Lys Gly His Asn Ala Ile Glu	
100 105 110	
ATG CAA ATG CAT TTA GAA TCA TTA TTA AGG ACT GAG TAT AGT ATG GAA	384
Met Gln Met His Leu Glu Ser Leu Leu Arg Thr Glu Tyr Ser Met Glu	
115 120 125	
CCG TGG ACA TTA CAA GAA ACA AGT TAT GAA ATG TGG CAA ACA CCA CCT	432
Pro Trp Thr Leu Gln Glu Thr Ser Tyr Glu Met Trp Gln Thr Pro Pro	
130 135 140	
AAA CGC TGT TTT AAA AAA CGG GGC AAA ACT GTA GAA GTT AAA TTT GAT	480
Lys Arg Cys Phe Lys Lys Arg Gly Lys Thr Val Glu Val Lys Phe Asp	

145		150		155		160	
GGC TGT GCA AAC AAT ACA ATG GAT TAT GTG GTA TGG ACA GAT GTG TAT	528						
Gly Cys Ala Asn Asn Thr Met Asp Tyr Val Val Trp Thr Asp Val Tyr							
		165		170		175	
GTG CAG GAC AAT GAC ACC TGG GTA AAG GTG CAT AGT ATG GTA GAT GCT	576						
Val Gln Asp Asn Asp Thr Trp Val Lys Val His Ser Met Val Asp Ala							
		180		185		190	
AAG GGT ATA TAT TAC ACA TGT GGA CAA TTT AAA ACA TAT TAT GTA AAC	624						
Lys Gly Ile Tyr Tyr Thr Cys Gly Gln Phe Lys Thr Tyr Tyr Val Asn							
		195		200		205	
TTT GTA AAA GAG GCA GAA AAG TAT GGG AGC ACC AAA CAT TGG GAA GTA	672						
Phe Val Lys Glu Ala Glu Lys Tyr Gly Ser Thr Lys His Trp Glu Val							
		210		215		220	
TGT TAT GGC AGC ACA GTT ATA TGT TCT CCT GCA TCT GTA TCT AGC ACT	720						
Cys Tyr Gly Ser Thr Val Ile Cys Ser Pro Ala Ser Val Ser Ser Thr							
		225		230		235	
ACA CAA GAA GTA TCC ATT CCT GAA TCT ACT ACA TAC ACC CCC GCA CAG	768						
Thr Gln Glu Val Ser Ile Pro Glu Ser Thr Tyr Thr Pro Ala Gln							
		245		250		255	
ACC TCC ACC CTT GTG TCC TCA AGC ACC AAG GAA GAC GCA GTG CAA ACG	816						
Thr Ser Thr Leu Val Ser Ser Ser Thr Lys Glu Asp Ala Val Gln Thr							
		260		265		270	
CCG CCT AGG AAA CGA GCA CGA GGA GTC CAA CAG TCC CCT TGC AAC GCC	864						
Pro Pro Arg Lys Arg Ala Arg Gly Val Gln Gln Ser Pro Cys Asn Ala							
		275		280		285	
TTG TGT GTG GCC CAC ATT GGA CCC GTG GAC AGT GGA AAC CAC AAC CTC	912						
Leu Cys Val Ala His Ile Gly Pro Val Asp Ser Gly Asn His Asn Leu							
		290		295		300	
ATC ACT AAC AAT CAC GAC CAG CAC CAA AGA CGG AAC AAC AGT AAC AGT	960						
Ile Thr Asn Asn His Asp Gln His Gln Arg Arg Asn Asn Ser Asn Ser							
		305		310		315	
TCA GCT ACG CCT ATA GTG CAA TTT CAA GGT GAA TCC AAT TGT TTA AAG	1008						
Ser Ala Thr Pro Ile Val Gln Phe Gln Gly Glu Ser Asn Cys Leu Lys							
		325		330		335	
TGT TTT AGA TAT AGG CTA AAT GAC AGA CAC AGA CAT TTA TTT GAT TTA	1056						
Cys Phe Arg Tyr Arg Leu Asn Asp Arg His Arg His Leu Phe Asp Leu							
		340		345		350	
ATA TCA TCA ACG TGG CAC TGG GCC TCC TCA AAG GCA CCA CAT AAA CAT	1104						
Ile Ser Ser Thr Trp His Trp Ala Ser Ser Lys Ala Pro His Lys His							
		355		360		365	
GCC ATT GTA ACT GTA ACA TAT GAT AGT GAG GAA CAA AGG CAA CAG TTT	1152						
Ala Ile Val Thr Val Thr Tyr Asp Ser Glu Glu Gln Arg Gln Gln Phe							
		370		375		380	
TTA GAT GTT GTA AAA ATA CCC CCT ACC ATT AGC CAC AAA CTG GGA TTT	1200						
Leu Asp Val Val Lys Ile Pro Pro Thr Ile Ser His Lys Leu Gly Phe							
		385		390		395	
ATG TCA CTG CAC CTA TTG GTC GAC ATG GGA GCA CCA AAC ATT GGG AAG	1248						
Met Ser Leu His Leu Leu Val Asp Met Gly Ala Pro Asn Ile Gly Lys							
		405		410		415	
TAT GTT ATG GCA GCA CAG TTA TAT GTT CTC CTG CAT CTG TAT CTA GCA	1296						
Tyr Val Met Ala Ala Gln Leu Tyr Val Leu Leu His Leu Tyr Leu Ala							
		420		425		430	
CTA CAC AAG AAG TAT CCA TTC CTG AAT CTA CTA CAT ACA CCC CCG CAC	1344						
Leu His Lys Lys Tyr Pro Phe Leu Asn Leu Leu His Thr Pro Pro His							
		435		440		445	

AGA Arg 450	CCT Pro 450	CCA Pro 450	CCC Pro 450	TTG Leu 450	TGT Cys 455	CCT Pro 455	CAA Gln 455	GCA Ala 455	CCA Pro 455	AGG Arg 460	AAG Lys 460	ACG Thr 460	CAG Gln 460	TGC Cys 460	AAA Lys 460	1392
CGC Arg 465	CGC Arg 465	CTA Leu 465	GGA Gly 465	AAC Asn 470	GAG Glu 470	CAC His 470	GAG Glu 475	GAG Glu 475	TCC Ser 475	AAC Asn 475	AGT Ser 475	CCC Pro 480	CTT Leu 480	GCA Ala 480	ACG Thr 480	1440
CCT Pro 485	TGT Cys 485	GTG Val 485	TGG Trp 485	CCC Pro 485	ACA Thr 485	TTG Leu 490	GAC Asp 490	CCG Pro 490	TGG Trp 490	ACA Thr 495	GTG Val 495	GAA Glu 495	ACC Thr 495	ACA Thr 495	ACC Thr 495	1488
TCA Ser 500	TCA Ser 500	CTA Leu 500	ACA Thr 500	ATC Ile 500	ACG Thr 505	ACC Thr 505	AGC Ser 505	ACC Thr 505	AAA Lys 505	GAC Asp 510	GGA Gly 510	ACA Thr 510	ACA Thr 510	GTA Val 510	ACA Thr 510	1536
GTT Val 515	CAG Gln 515	CTA Leu 515	CGC Arg 515	CTA Leu 520	AGA Arg 520	TCT Ser 520	ATG Met 520	GAA Glu 520	GTG Val 525	GTG Val 525	CCT Pro 525	GTA Val 525	CAA Gln 525	ATA Ile 525	GCT Ala 525	1584
GCA Ala 530	GGA Gly 530	ACA Thr 530	ACC Thr 535	AGC Ser 535	ACA Thr 535	TTC Phe 535	ATA Ile 540	CTG Leu 540	CCT Pro 540	GTT Val 540	ATA Ile 540	ATT Ile 540	GCA Ala 540	TTT Phe 540	GTT Val 540	1632
GTA Val 545	TGT Cys 545	TTT Phe 545	GTT Val 550	AGC Ser 550	ATC Ile 550	ATA Ile 555	CTT Leu 555	ATT Ile 555	GTA Val 555	TGG Trp 555	ATA Ile 560	TCT Ser 560	GAG Glu 560	TTT Phe 560	ATT Ile 560	1680
GTG Val 565	TAC Tyr 565	ACA Thr 565	TCT Ser 570	GTG Val 570	CTA Leu 570	GTA Val 575	CTA Leu 575	ACA Thr 575	CTG Leu 580	CTT Leu 580	TTA Leu 585	TAT Tyr 585	TTA Leu 585	CTA Leu 590	TTG Leu 590	1728
TGG Trp 580	CTG Leu 580	CTA Leu 580	TTA Leu 580	ACA Thr 585	ACC Thr 585	CCC Pro 585	TTG Leu 585	CAA Gln 585	TTT Phe 590	TTC Phe 590	CTA Leu 590	CTA Leu 590	ACT Thr 590	CTA Leu 590	CTT Leu 590	1776
GTG Val 595	TGT Cys 595	TAC Tyr 595	TGT Cys 600	CCC Pro 600	GCA Ala 600	TTG Leu 600	TAT Tyr 605	ATA Ile 605	CAC His 605	TAC Tyr 605	TAT Tyr 605	ATT Ile 605	GTT Val 610	ACC Thr 610	ACA Thr 610	1824
CAG Gln 610	CAA Gln 610	CTT Leu 615	AAG Lys 615	ATG Met 615	ATG Met 615	CTA Leu 615	ACA Thr 620	TGT Cys 620	CAA Gln 620	TTT Phe 620	AAT Asn 620	GAT Asp 620	GGA Gly 620	GAT Asp 620	ACC Thr 620	1872
TGG Trp 625	CTG Leu 625	GGT Gly 630	TTG Leu 630	TGG Trp 630	TTG Leu 630	TTA Leu 635	TGT Cys 635	GCC Ala 635	TTT Phe 635	ATT Ile 640	GTA Val 640	GGG Gly 640	ATG Met 640	TTG Leu 640	GGG Gly 640	1920
TTA Leu 645	TTA Leu 645	TTG Leu 645	ATG Met 645	CAC His 645	TAT Tyr 645	AGA Arg 650	GCT Ala 650	GTA Val 650	CAA Gln 650	GGG Gly 655	GAT Asp 655	AAA Lys 655	CAC His 655	ACC Thr 655	AAA Lys 655	1968
TGT Cys 660	AAG Lys 660	AAG Lys 660	TGT Cys 660	AAC Asn 660	AAA Lys 665	CAC His 665	AAC Asn 665	TGT Cys 665	AAT Asn 665	GAT Asp 670	GAT Asp 670	TAT Tyr 670	GTA Val 670	ACT Thr 670	ATG Met 670	2016
CAT His 675	TAT Tyr 675	ACT Thr 675	ACT Thr 675	GAT Asp 680	GGT Gly 680	GAT Asp 680	TAT Tyr 680	ATA Ile 685	TAT Tyr 685	ATG Met 685	AAT Asn 685	CCG Pro 685	CGG Arg 685	ATG Met 685	GAA Glu 685	2064
AGT Ser 690	GCA Ala 690	AAT Asn 690	GCC Ala 695	TCC Ser 695	ACG Thr 695	TCT Ser 695	GCA Ala 695	ACG Thr 700	ACC Thr 700	ATA Ile 700	GAC Asp 700	CAG Gln 700	TTG Leu 700	TGC Cys 700	AAG Lys 700	2112
ACG Thr 705	TTT Phe 705	AAT Asn 710	CTA Leu 710	TCT Ser 710	ATG Met 710	CAT His 715	ACG Thr 715	TTG Leu 715	CAA Gln 715	ATT Ile 715	AAT Asn 720	TGT Cys 720	GTG Val 720	TTT Phe 720	TGC Cys 720	2160
AAG Lys 725	AAT Asn 725	GCA Ala 725	CTG Leu 725	ACC Thr 725	ACA Thr 730	GCA Ala 730	GAG Glu 730	ATT Ile 730	TAT Tyr 730	TCA Ser 735	TAT Tyr 735	GCA Ala 735	TAT Tyr 735	AAA Lys 735	CAC His 735	2208
CTA Leu 725	AAG Lys 725	GTC Val 725	CTG Leu 725	TTT Phe 725	CGA Arg 725	GGC Gly 725	GGC Gly 725	TAT Tyr 725	CCA Pro 725	TAT Tyr 725	GCA Ala 725	GCC Ala 725	TGC Cys 725	GCG Ala 725	TGC Cys 725	2256

740	745	750	
TGC CTA GAA TTT CAT GGA AAA ATA AAC CAA TAT AGA CAC TTT GAT TAT Cys Leu Glu Phe His Gly Lys Ile Asn Gln Tyr Arg His Phe Asp Tyr 755 760 765			2304
GCT GGA TAT GCA ACA ACA GTT GAA GAA GAA ACT AAA CAA GAC ATC TTA Ala Gly Tyr Ala Thr Thr Val Glu Glu Glu Thr Lys Gln Asp Ile Leu 770 775 780			2352
GAC GTG CTA ATT CGG TGC TAC CTG TGT CAC AAA CCG CTG TGT GAA GTA Asp Val Leu Ile Arg Cys Tyr Leu Cys His Lys Pro Leu Cys Glu Val 785 790 795 800			2400
GAA AAG GTA AAA CAT ATA CTA ACC AAG GCG CGG TTC ATA AAG CTA AAT Glu Lys Val Lys His Ile Leu Thr Lys Ala Arg Phe Ile Lys Leu Asn 805 810 815			2448
TGT ACG TGG AAG GGT CGC TGC CTA CAC TGC TGG ACA ACA TGC ATG GAA Cys Thr Trp Lys Gly Arg Cys Leu His Cys Trp Thr Thr Cys Met Glu 820 825 830			2496
GAC ATG TTA CCC GCT AGC ATG CAT GGA AGA CAT GTT ACC CTA AAG GAT Asp Met Leu Pro Ala Ser Met His Gly Arg His Val Thr Leu Lys Asp 835 840 845			2544
ATT GTA TTA GAC CTG CAA CCT CCA GAC CCT GTA GGG TTA CAT TGC TAT Ile Val Leu Asp Leu Gln Pro Pro Asp Pro Val Gly Leu His Cys Tyr 850 855 860			2592
GAG CAA TTA GTA GAC AGC TCA GAA GAT GAG GTG GAC GAA GTG GAC GGA Glu Gln Leu Val Asp Ser Ser Glu Asp Glu Val Asp Glu Val Asp Gly 865 870 875 880			2640
CAA GAT TCA CAA CCT TTA AAA CAA CAT TTC CAA ATA GTG ACC TGT TGC Gln Asp Ser Gln Pro Leu Lys Gln His Phe Gln Ile Val Thr Cys Cys 885 890 895			2688
TGT GGA TGT GAC AGC AAC GTT CGA CTG GTT GTG CAG TGT ACA GAA ACA Cys Gly Cys Asp Ser Asn Val Arg Leu Val Val Gln Cys Thr Glu Thr 900 905 910			2736
GAC ATC AGA GAA GTG CAA CAG CTT CTG TTG GGA ACA CTA AAC ATA GTG Asp Ile Arg Glu Val Gln Gln Leu Leu Leu Gly Thr Leu Asn Ile Val 915 920 925			2784
TGT CCC ATC TGC GCA CCG AAG ACC CGA TCG ATG GCG GAC GAT TCA GGT Cys Pro Ile Cys Ala Pro Lys Thr Arg Ser Met Ala Asp Asp Ser Gly 930 935 940			2832
ACA GAA AAT GAG GGG TCT GGG TGT ACA GGA TGG TTT ATG GTA GAA GCT Thr Glu Asn Glu Gly Ser Gly Cys Thr Gly Trp Phe Met Val Glu Ala 945 950 955 960			2880
ATA GTG CAA CAC CCA ACA GGT ACA CAA ATA TCA GAC GAT GAG GAT GAG Ile Val Gln His Pro Thr Gly Thr Gln Ile Ser Asp Asp Glu Asp Glu 965 970 975			2928
GAG GTG GAG GAC AGT GGG TAT GAC ATG GTG GAC TTT ATT GAT GAC AGC Glu Val Glu Asp Ser Gly Tyr Asp Met Val Asp Phe Ile Asp Asp Ser 980 985 990			2976
AAT ATT ACA CAC AAT TCA CTG GAA GCA CAG GCA TTG TTT AAC AGG CAG Asn Ile Thr His Asn Ser Leu Glu Ala Gln Ala Leu Phe Asn Arg Gln 995 1000 1005			3024
GAG GCG GAC ACC CAT TAT GCG ACT GTG CAG GAC CTA AAA CGA AAG TAT Glu Ala Asp Thr His Tyr Ala Thr Val Gln Asp Leu Lys Arg Lys Tyr 1010 1015 1020			3072
TTA GGT AGT CCA TAT GTT AGT CCT ATA AAC ACT ATA GCC GAG GCA GTG Leu Gly Ser Pro Tyr Val Ser Pro Ile Asn Thr Ile Ala Glu Ala Val 1025 1030 1035 1040			3120

GAA AGT GAA ATA AGT CCA CGA TTG GAC GCC ATT AAA CTT ACA AGA CAG 3168
 Glu Ser Glu Ile Ser Pro Arg Leu Asp Ala Ile Lys Leu Thr Arg Gln
 1045 1050 1055

CCA AAA AAG GTA AAG CGA CGG CTG TTT CAA ACC AGG GAA CTA ACG GAC 3216
 Pro Lys Lys Val Lys Arg Arg Leu Phe Gln Thr Arg Glu Leu Thr Asp
 1060 1065 1070

AGT GGA TAT GGC TAT TCT GAA GTG GAA GCT GGA ACG GGA ACG CAG GTA 3264
 Ser Gly Tyr Gly Tyr Ser Glu Val Glu Ala Gly Thr Gly Thr Gln Val
 1075 1080 1085

GAG AAA CAT GGC GTA CCG GAA AAT GGG GGA GAT GGT CAG GAA AAG GAC 3312
 Glu Lys His Gly Val Pro Glu Asn Gly Gly Asp Gly Gln Glu Lys Asp
 1090 1095 1100

ACA GGA AGG GAC ATA GAG GGG GAG GAA CAT ACA GAG GCG GAA GCG CCC 3360
 Thr Gly Arg Asp Ile Glu Gly Glu Glu His Thr Glu Ala Glu Ala Pro
 1105 1110 1115 1120

ACA AAC AGT GTA CGG GAG CAT GCA GGC ACA GCA GGA ATA TTG GAA TTG 3408
 Thr Asn Ser Val Arg Glu His Ala Gly Thr Ala Gly Ile Leu Glu Leu
 1125 1130 1135

TTA AAA TGT AAA GAT TTA CGG GCA GCA TTA CTT GGT AAG TTT AAA GAA 3456
 Leu Lys Cys Lys Asp Leu Arg Ala Ala Leu Leu Gly Lys Phe Lys Glu
 1140 1145 1150

TGC TTT GGG CTG TCT TTT ATA GAT TTA ATT AGG CCA TTT AAA AGT GAT 3504
 Cys Phe Gly Leu Ser Phe Ile Asp Leu Ile Arg Pro Phe Lys Ser Asp
 1155 1160 1165

AAA ACA ACA TGT TTA GAT TGG GTG GTA GCA GGG TTT GGT ATA CAT CAT 3552
 Lys Thr Thr Cys Leu Asp Trp Val Val Ala Gly Phe Gly Ile His His
 1170 1175 1180

AGC ATA TCA GAG GCA TTT CAA AAA TTA ATT GAG CCA TTA AGT TTA TAT 3600
 Ser Ile Ser Glu Ala Phe Gln Lys Leu Ile Glu Pro Leu Ser Leu Tyr
 1185 1190 1195 1200

GCA CAT ATA CAA TGG CTA ACA AAT GCA TGG GGA ATG GTA TTG TTA GTA 3648
 Ala His Ile Gln Trp Leu Thr Asn Ala Trp Gly Met Val Leu Leu Val
 1205 1210 1215

TTA TTA AGA TTT AAA GTA AAT AAA AGT AGA AGT ACC GTT GCA CGT ACA 3696
 Leu Leu Arg Phe Lys Val Asn Lys Ser Arg Ser Thr Val Ala Arg Thr
 1220 1225 1230

CTT GCA ACG CTA TTA AAT ATA CCT GAA AAC CAA ATG TTA ATA GAG CCA 3744
 Leu Ala Thr Leu Leu Asn Ile Pro Glu Asn Gln Met Leu Ile Glu Pro
 1235 1240 1245

CCA AAA ATA CAA AGT GGT GTT GCA GCC CTG TAT TGG TTT CGT ACA GGT 3792
 Pro Lys Ile Gln Ser Gly Val Ala Ala Leu Tyr Trp Phe Arg Thr Gly
 1250 1255 1260

ATA TCA AAT GCC AGT ACA GTT ATA GGG GAA GCA CCA GAA TGG ATA ACA 3840
 Ile Ser Asn Ala Ser Thr Val Ile Gly Glu Ala Pro Glu Trp Ile Thr
 1265 1270 1275 1280

CGC CAA ACA GTT ATT GAA CAC GGG TTG GCA GAC AGT CAG TTT AAA TTA 3888
 Arg Gln Thr Val Ile Glu His Gly Leu Ala Asp Ser Gln Phe Lys Leu
 1285 1290 1295

ACA GAA ATG GTG CAG TGG GCG TAT GAT AAT GAC ATA TGC GAG GAG AGT 3936
 Thr Glu Met Val Gln Trp Ala Tyr Asp Asn Asp Ile Cys Glu Glu Ser
 1300 1305 1310

GAA ATT GCA TTT GAA TAT GCA CAA AGG GGA GAT TTT GAT TCT AAT GCA 3984
 Glu Ile Ala Phe Glu Tyr Ala Gln Arg Gly Asp Phe Asp Ser Asn Ala
 1315 1320 1325

CGA GCA TTT TTA AAT AGC AAT ATG CAG GCA AAA TAT GTG AAA GAT TGT 4032
 Arg Ala Phe Leu Asn Ser Asn Met Gln Ala Lys Tyr Val Lys Asp Cys
 1330 1335 1340

1330	1335	1340	
GCA ACT ATG TGT AGA CAT TAT AAA CAT GCA GAA ATG AGG AAG ATG TCT Ala Thr Met Cys Arg His Tyr Lys His Ala Glu Met Arg Lys Met Ser 1345 1350 1355 1360			4080
ATA AAA CAA TGG ATA AAA CAT AGG GGT TCT AAA ATA GAA GGC ACA GGA Ile Lys Gln Trp Ile Lys His Arg Gly Ser Lys Ile Glu Gly Thr Gly 1365 1370 1375			4128
AAT TGG AAA CCA ATT GTA CAA TTC CTA CGA CAT CAA AAT ATA GAA TTC Asn Trp Lys Pro Ile Val Gln Phe Leu Arg His Gln Asn Ile Glu Phe 1380 1385 1390			4176
ATT CCT TTT TTA ACT AAA TTT AAA TTA TGG CTG CAC GGT ACG CCA AAA Ile Pro Phe Leu Thr Lys Phe Lys Leu Trp Leu His Gly Thr Pro Lys 1395 1400 1405			4224
AAA AAC TGC ATA GCC ATA GTA GGC CCT CCA GAT ACT GGG AAA TCG TAC Lys Asn Cys Ile Ala Ile Val Gly Pro Pro Asp Thr Gly Lys Ser Tyr 1410 1415 1420			4272
TTT TGT ATG AGT TTA ATA AGC TTT CTA GGA GGT ACA GTT ATT AGT CAT Phe Cys Met Ser Leu Ile Ser Phe Leu Gly Gly Thr Val Ile Ser His 1425 1430 1435 1440			4320
GTA AAT TCC AGC AGC CAT TTT TGG TTG CAA CCG TTA GTA GAT GCT AAG Val Asn Ser Ser Ser His Phe Trp Leu Gln Pro Leu Val Asp Ala Lys 1445 1450 1455			4368
GTA GCA TTG TTA GAT GAT GCA ACA CAG CCA TGT TGG ATA TAT ATG GAT Val Ala Leu Leu Asp Asp Ala Thr Gln Pro Cys Trp Ile Tyr Met Asp 1460 1465 1470			4416
ACA TAT ATG AGA AAT TTG TTA GAT GGT AAT CCT ATG AGT ATT GAC AGA Thr Tyr Met Arg Asn Leu Leu Asp Gly Asn Pro Met Ser Ile Asp Arg 1475 1480 1485			4464
AAG CAT AAA GCA TTG ACA TTA ATT AAA TGT CCA CCT CTG CTA GTA ACG Lys His Lys Ala Leu Thr Leu Ile Lys Cys Pro Pro Leu Leu Val Thr 1490 1495 1500			4512
TCC AAC ATA GAT ATT ACT AAA GAA GAT AAA TAT AAG TAT TTA CAT ACT Ser Asn Ile Asp Ile Thr Lys Glu Asp Lys Tyr Lys Tyr Leu His Thr 1505 1510 1515 1520			4560
AGA GTA ACA ACA TTT ACA TTT CCA AAT CCA TTC CCT TTT GAC AGA AAT Arg Val Thr Thr Phe Thr Phe Pro Asn Pro Phe Pro Phe Asp Arg Asn 1525 1530 1535			4608
GGG AAT GCA GTG TAT GAA CTG TCA AAT ACA AAC TGG AAA TGT TTT TTT Gly Asn Ala Val Tyr Glu Leu Ser Asn Thr Asn Trp Lys Cys Phe Phe 1540 1545 1550			4656
GAA AGA CTG TCG TCA AGC CTA GAC ATT CAG GAT TCT GAG GAC GAG GAA Glu Arg Leu Ser Ser Ser Leu Asp Ile Gln Asp Ser Glu Asp Glu Glu 1555 1560 1565			4704
GAT GGA AGC AAT AGC CAA GCG TTT AGA TGC GTG CCA GGA ACA GTT GTT Asp Gly Ser Asn Ser Gln Ala Phe Arg Cys Val Pro Gly Thr Val Val 1570 1575 1580			4752
AGA ACT TTA TGAGGTACC Arg Thr Leu 1585			4770

CLAIMS:

1. A polyprotein construct comprising at least two amino acid sequences fused directly or indirectly together, each of said sequences being the sequence of an early ORF protein of papillomavirus (PV) or an immunogenic variant or fragment thereof, and at least one of said sequences being other than the E6 or E7 protein sequence or an immunogenic variant or fragment thereof.
2. A polyprotein construct according to claim 1, wherein said sequences are sequences of early ORF proteins of human PV, or immunogenic variants or fragments thereof.
3. A polyprotein construct according to claim 2, wherein said early ORF proteins are selected from the group consisting of the E1, E2, E3, E4, E5 (E5a, E5b), E6, E7 and E8 proteins of PV.
4. A polyprotein construct according to any of claims 1 to 3, selected from the group consisting of:
 - (a) E6/E4
 - (b) E6/E5a/E4
 - (c) E6/E7/E4
 - (d) E6/E7/E5a/E4
 - (e) E6/E7/E1/E4
 - (f) E6/E7/E5a/E1/E4
 - (g) E6/E7/E5a/E1/E2/E4
 - (h) E6/E7/E5a/E5b/E1/E2/E4
 - (i) E2/E5b
 - (j) E2/E1/E5b
 - (k) E2/E5a/E5b
 - (l) E2/E1/E5a/E5b

- 42 -

- (m) E2/E4/E5a/E5b/E6/E7/E1
- (n) E2/E3/E4/E5/E8/E6/E7/E1.


5. A polyprotein construct according to claim 1, further comprising one or more linker sequences between and/or before and/or after said amino acid sequences.
6. A polyprotein construct according to claim 5, wherein said linker sequence(s) comprise from 1 to 5 amino acid residues.
7. A polyprotein construct according to claim 1, further comprising a tag protein or peptide moiety fused or otherwise coupled thereto.
8. A polyprotein construct according to claim 7, wherein said tag moiety is selected from the group consisting of (his)₆, glutathione-S-transferase (GST) and FLAG.
9. A polyprotein construct according to claim 1, further comprising an adjuvant moiety fused or otherwise coupled thereto.
10. A polyprotein construct according to claim 9, wherein said adjuvant moiety is selected from diphtheria toxin, cholera toxin and *E. coli* heat labile toxin (LT) and non-toxic derivatives thereof such as the holotoxoid or B sub-unit of cholera toxin or LT.
11. A polyprotein construct according to claim 1, further comprising a lipid binding region.
12. A polyprotein construct according to claim 11, wherein said lipid binding region is an influenza haemagglutinin tail.

- 43 -

13. A composition for eliciting a humoral and/or cellular immune response against papillomavirus in a host animal, said composition comprising an immunologically effective amount of a polyprotein construct according to any of claims 1 to 12, together with a pharmaceutically acceptable carrier and/or diluent.
14. A vaccine composition according to claim 13, further comprising an adjuvant.
15. A method for eliciting a humoral and/or cellular response against papillomavirus in a host animal, which method comprises administering to the host animal an immunologically effective amount of a polyprotein construct according to any of claims 1 to 12.
16. A method according to claim 15, wherein said polyprotein construct is administered in a composition together with a pharmaceutically acceptable carrier and/or diluent.
17. A method according to claim 16, wherein said composition further comprises an adjuvant.
18. A method according to any of claims 15 to 17, wherein said host animal is a human.
19. Use of a polyprotein construct according to any of claims 1 to 12, in eliciting an immune response against papillomavirus in a host animal.
20. A nucleic acid molecule which encodes a polyprotein construct according to any of claims 1 to 12.
21. A recombinant DNA molecule comprising an expression control sequence operatively linked to a nucleic acid molecule according to claim 20.

22. A recombinant DNA molecule according to claim 21, wherein said expression control sequence comprises promoter and initiator sequences, the sequence of nucleotides encoding the polyprotein construct being located in a single translational frame 3' to the promoter and initiator sequences, and a termination sequence located 3' to said sequence of nucleotides.
23. A recombinant DNA cloning vehicle or vector comprising a recombinant DNA molecule according to claim 21 or claim 22.
24. A recombinant DNA cloning vehicle or vector according to claim 23, wherein said vector is a plasmid.
25. A host cell transfected or transformed with a recombinant DNA molecule according to claim 21 or claim 22, or a recombinant DNA cloning vehicle or vector according to claim 23 or claim 24.
26. A host cell according to claim 25, wherein said host cell is *E. coli*.
27. A recombinant polyprotein construct prepared by expression in a host cell according to claim 25 or claim 26.
28. A composition comprising a nucleic acid molecule according to claim 20, together with a pharmaceutically acceptable carrier and/or diluent.
29. A method for eliciting an immune response against PV in a host animal, which method comprises administering to the host animal an immunologically effective amount of a nucleic acid molecule according to claim 20.
30. Use of a nucleic acid molecule according to claim 20 in eliciting an immune response against PV in a host animal.

Figure 1



2/12





FIGURE 2

SUBSTITUTE SHEET (RULE 26)

3/12

Figure 3



SUBSTITUTE SHEET (RULE 26)

4/12

ANTI-HEXAHIS (DIANOVA) Mab




Figure 4.

SUBSTITUTE SHEET (RULE 26)

5/12

ANTI-E4 PEPTIDE





Figure 5.

SUBSTITUTE SHEET (RULE 26)

6/12


Figure 6



SUBSTITUTE SHEET (RULE 26)

7/12


Figure 7



SUBSTITUTE SHEET (RULE 26)

8/12


Figure 8



SUBSTITUTE SHEET (RULE 26)

9/12


Figure 9



SUBSTITUTE SHEET (RULE 26)


10/12

Figure 10



11/12


Figure 11



SUBSTITUTE SHEET (RULE 26)

12/12

Figure 12



100 aa

 hexaHis Tag encoded by pTrcHisA

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/AU 96/00473

A. CLASSIFICATION OF SUBJECT MATTER

Int Cl⁶: C07K 14/025; C12N 15/37, 15/86, 5/10; A61K 39/12, 31/73

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC⁶: C07K, C12N, A61K. Chemical Abstracts. All through Electronic Databases

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
DERWENT Databases: WPAT & JAPIO. Search terms: See extra sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	DE 4435907 (GUTZMANN et al), 11 April 1996, IPC ⁶ C07K 14/37, 14/01, 14/08; A61K 38/16 See claims, especially claims 9 and 10	1-3
X	TANIGUCHI & YASUMOTO: "A Major Transcript of Human Papillomavirus Type 16 in Transformed NIH 3T3 Cells contains Polycistronic mRNA encoding E7, E5, and E1^E4 Fusion Gene". Virus Genes, 3(3), pp 221-233, 1990. See abstract, figures 3 and 6, p 229 lines 4-10 and 15-16	1-3

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search
10 September 1996

Date of mailing of the international search report
18.09.96

Name and mailing address of the ISA/AU
AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION
PO BOX 200
WODEN ACT 2606
AUSTRALIA Facsimile No.: (06) 285 3929

Authorized officer
ROBYN PORTER
Telephone No.: (06) 283 2318

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/AU 96/00473

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROHLFS et al: "Viral Transcription in Human Keratinocyte Cell Lines Immortalized by Human Papillomavirus Type-16". Virology, 183, pp 331-342 (1991). See Figure 1; page 334, column 2, lines 5-9, 13-15 and 19-20; page 335 column 1, lines 26-27 and column 2 lines 2-10	1-4
X	CHIANG et al: "An E1M ⁺ E2C Fusion Protein Encoded by Human Papillomavirus Type 11 Is a Sequence-Specific Transcription Repressor". Journal of Virology, 65(6), pp 3317-3329, 1991. See abstract, p 3318, column 2, 2nd full paragraph, Figures 1 and 2, p 3321, column 1, 1st full sentence, column 2, line 3 - p 1322, column 1, line 2, column 2 lines 2-5, p 3323, column 1, 1st full paragraph, p 3326, column 2 lines 5-9 and 1st 2 sentences of 1st full paragraph	1-4, 20-22
X	LAMBERTI et al: "Transcriptional activation by the papillomavirus E6 zinc finger oncoprotein". The EMBO Journal, 9 (6), pp 1907-1913, 1990. See abstract, figure 1, p 1912, 2nd paragraph of "Constructions"	1, 5, 20-22
X	WO 92/11290 (CETUS CORPORATION), 9 July 1992, IPC ⁵ C07K 13/00, 15/18; A61K 37/10; G01N 33/569, 33/68; C12Q 1/18, 1/70. See abstract, p 6 lines 3-11, page 7 line 12 - page 12, line 9, claims	1-3, 13-20
X	TOMITA & SIMIZU: "Translational properties of the human papillomavirus type-6 L1-coding mRNA". Gene, 133, pp 223-225, 1993. See in particular figure 1B #3	1-3, 5, 20, 21
X	WO 94/12629 (BAYLOR COLLEGE OF MEDICINE) 9 June 1994, IPC ⁵ C12N 15/00; A61K 31/70. See abstract, p 3 lines 16-31, p 7 line 19 - p8 line 7, p11 lines 1-15, p 16 lines 35-36, Example 1 (on p 27), claim 1, Figure 1	1-3, 5, 20, 21

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/AU 96/00473

Box

Search terms used:

WPAT and JAPIO search

SS1 : PAPILLOMAVIRUS## OR PAPILLOMA(W)VIRUS##

SS2 : EARLY (3N)(ORF OR OPEN(W)READING(W)FRAME# OR PROTEIN# OR POLYPEPTIDE#)

SS3 : 1 AND 2

SS4 : E1# OR E2# OR E3# OR E4# OR E5# OR E6# OR E7# OR E8#

SS5 : 1 AND 4

SS 6 : 3 OR 5

Search terms used:

Chemical Abstracts Search

L1 : S EARLY (3N) (ORF OR OPEN()READING()FRAME# OR PROTEIN# OR POLYPEPTIDE#)/IT

L2 : S PAPILLOMAVIRUS?/IT OR PAPILLOMA()VIRUS##/IT

L3 : S (E1# OR E2# OR E3# OR E4# OR E5# OR E6# OR E7# OR E8#)/IT

L4 : S L1 AND L2

L5 : S L3 (L) L2

L6 : S (FUS## OR FUSI##)/IT

L7 : S L6 (L) L5

L8 : S L6(L) L4

L9 : S L7 OR L8

International Application No.
PCT/AU 96/00473

Patent Document Cited in Search Report				Patent Family Member			
DE	4435907	AU	42701/96	WO	9611272		
WO	9211290	AU JP	91731/91 7503230	CA US	2098926 5464936	EP	563307
WO	9412629	AU	60140/94				

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)